arXiv:1103.0380v1 [math.AG] 2 Mar 2011

A CONSTRUCTIVE APPROACH TO THE CLASSIFICATION OF WONDERFUL VARIETIES

P. BRAVI AND G. PEZZINI

Abstract. Let G be an adjoint semisimple linear group over $\mathbb{C}$. We analyze the relationship between the algebraic and the combinatorial point of view on Luna's theory of wonderful G-varieties. Some techniques relating a spherical system to its associated wonderful subgroup are discussed; as a byproduct, we reduce for any group G the proof of the classification to its verification on a small family of spherical systems, called primitive.

1. Introduction

Luna's theory of wonderful varieties establishes a connection between *wonderful G-varieties* for a semisimple complex algebraic group G, and certain combinatorial objects called *spherical systems* of G ([14]). The main conjecture, i.e. that spherical systems classify these varieties, is a work in progress and has been verified in many particular cases ([14, 15, 5, 1, 3]). Uniqueness of the wonderful variety having a given spherical system has been proved by Losev in [12], and a proof of the whole conjecture has been proposed by Cupit-Foutou in [7] and [8] using a different method.

Beyond the classification, the theory describes many geometric constructions and their combinatorial counterparts. For example, morphisms between varieties correspond to *quotients* of spherical systems.

In this article we develop some techniques relating a spherical system to the generic stabilizer (called a *wonderful subgroup* of G) of the corresponding variety. Whenever such a wonderful variety exists, we say that the spherical system is *geometrically realizable*.

Two approaches to these techniques are possible. The first one is described in §3, and extracts informations on the wonderful subgroup H only from the spherical system $\mathscr{S}$ itself. This approach is direct, but yet unable to provide a complete description of H, nor of a candidate H in cases where geometric realizability of $\mathscr{S}$ is not assured.

With this point of view, a Levi part of H (as already pointed out in [12]), can be guessed using the classification of reductive spherical subgroups of semisimple groups and the knowledge of their spherical systems. The unipotent radical is more elusive, although some strategy for it seems to be possible. Our study of H is also in some sense non-canonical, since it requires the choice of a parabolic subgroup of G minimal containing H: examples show that such a choice is not unique.

The second approach, developed in §§4–6, involves the study of particular quotients of $\mathscr{S}$, which are supposed to be geometrically realizable. Here we can produce a precise description for H, proving at the same time the geometric realizability of

$\mathscr{S}$. However, all is done assuming existence and knowledge of the generic stabilizers of some quotient of $\mathscr{S}$.

We also review some results that were already known, always about the relationship between spherical systems and wonderful subgroups. We refine the results of [14] on fiber products of wonderful varieties, and discuss the so-called projective fibrations (see [14, §3.6]) without restrictions on G as a particular case of a more general situation: *quotients of higher defect*.

All our work leads to the definition of *primitive* spherical systems, a direct generalization of the notion defined in [14] and subsequent works.

The corresponding wonderful varieties, called again *primitive*, play a significative role in the theory. In an analogy with representation theory they would correspond to indecomposable representations. In addition, the smaller class of primitive varieties admitting no quotient of higher defect would correspond to simple representations.

Primitive spherical systems can also be used to provide a proof of the existence part of Luna's conjecture, in the same spirit of [14]. Namely, we prove that Luna's conjecture follows from case-by-case analysis on primitive spherical systems without quotients of higher defect. This will be the subject of forthcoming papers.

Acknowledgments. We are grateful to Prof. D. Luna for his comments and suggestions and for pointing out some errors in a previous version of this paper, and to Prof. F. Knop for useful discussions. The secondly-named author was supported by the DFG Schwerpunktprogramm 1388 – Darstellungstheorie.

2. Definitions

2.1. **Wonderful varieties.**

Throughout this work, G will be a semisimple complex algebraic group, T a maximal torus, B a Borel subgroup containing T, S the corresponding set of simple roots of the root system of (G,T).

For all $S'\subset S$, $P_{S'}$ denotes the corresponding parabolic subgroup containing B (namely, S' is a set of simple roots of the root system of (L,T), where $L=L_{S'}$ is the Levi subgroup of $P_{S'}$ containing T). Similarly, $P_{-S'}$ denotes the opposite parabolic subgroup of $P_{S'}$, with respect to the maximal torus T. We also denote by $G_{S'}$ the semisimple part $(L_{S'},L_{S'})$ of $L_{S'}$.

For any parabolic subgroup P containing T, L_P will be its "standard" Levi subgroup containing T.

If H is any affine algebraic group, $\mathrm{Lie}\,H$ will denote its Lie algebra and H° its connected component containing the unit element $e\in H$, H^r will be its radical and H^u its unipotent radical, $Z(H)$ will be its center, $\mathrm{N}_K(H)$ will be its normalizer in K.

We are now going to recall the basic definitions about wonderful varieties, see [14] and [4] for further details and references.

Definition 2.1.1. A G-variety is called *wonderful* of rank r if and only if it is smooth, complete, with r smooth prime G-divisors $D_1,\ldots,D_r$ with normal crossings, such that the G-orbit closures are exactly all the intersections $\cap_{i\in I}D_i$, for any $I\subseteq\{1,\ldots,r\}$.

We will always assume that $Z(G)$ acts trivially on a wonderful G-variety. This is justified by the results of Luna in [14], where the classification is reduced to the case where G is adjoint.

A wonderful G-variety is known to be projective and spherical.

Let X be a wonderful G-variety. There exists a unique point $z \in X$ stabilized by B_- (where B_- is the Borel subgroup opposite to B with respect to T, i.e. $B_- \cap B = T$). The orbit $G.z$ is the unique closed orbit, $\cap_{i=1}^r D_i$; the parabolic subgroup opposite to G_z, usually denoted by P_X, is the stabilizer of the open B-orbit of X. Set S^p_X such that $P_X = P_{S^p_X}$.

The T-weights occurring in the normal space $T_zX/T_zG.z$ of $G.z$ at z are called spherical roots, their set is denoted by Σ_X. Spherical roots are in bijective correspondence with prime G-divisors, say $\sigma \mapsto D^\sigma$, and form a basis of the lattice of B-weights in $\mathbb{C}(X)$ (we are identifying B-weights with T-weights).

The not G-stable prime B-divisors of X are called colors, their set is denoted by Δ_X. The colors are representatives of a basis of Pic$X$, so one can define a $\mathbb{Z}$-bilinear pairing, called Cartan pairing, $c_X\colon \mathbb{Z}\Delta_X \times \mathbb{Z}\Sigma_X \to \mathbb{Z}$ such that $[D^\sigma] = \sum_{D\in\Delta_X} c_X(D,\sigma)[D]$, for all $\sigma \in \Sigma_X$.

This is also denoted as a map $\rho_X\colon \Delta_X \to \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}\Sigma_X, \mathbb{Q})$, where $\rho_X(D)(-) = c_X(D,-)$.

For all $\alpha \in S$, set $\Delta_X(\alpha) = \{D \in \Delta_X : P_{\{\alpha\}}.D \neq D\}$. One has $\cup_{\alpha\in S}\Delta_X(\alpha) = \Delta_X$ and, for all $\alpha \in S$, $\mathrm{card}(\Delta_X(\alpha)) \leq 2$. Clearly, $\alpha \in S^p_X$ if and only if $\Delta_X(\alpha) = \emptyset$. Moreover, if $D \in \Delta_X(\alpha)$ and $\mathrm{card}(\Delta_X(\alpha)) = 1$ then $\rho_X(D)$ is uniquely determined by α. One has $\mathrm{card}(\Delta_X(\alpha)) = 2$ if and only if $\alpha \in S \cap \Sigma_X$, in this case, say $\Delta_X(\alpha) = \{D^+, D^-\}$, $\rho_X(D^+)$ and $\rho_X(D^-)$ are not always determined by α, but their sum is. Let $\mathbf{A}_X$ denote the subset of colors $D \in \cup_{\alpha\in S\cap\Sigma_X}\Delta_X(\alpha)$ endowed with the $\mathbb{Z}$-linear functionals $\rho_X(D)$.

Definition 2.1.2. The datum of $(S^p_X, \Sigma_X, \mathbf{A}_X)$, also denoted by $\mathscr{S}_X$, is called the spherical system of X.

Any G-orbit closure X' of X is a wonderful G-variety itself. Its set of spherical roots $\Sigma' = \Sigma_{X'}$ is a subset of Σ_X, and X' is called the localization of X with respect to Σ' and denoted by $X' = X_{\Sigma'}$. Indeed, $X_{\Sigma'} = \cap_{\sigma\in\Sigma_X\setminus\Sigma'} D^\sigma$. The spherical system $\mathscr{S}_{X'}$ of X' is given by $S^p_{X'} = S^p_X$, $\Sigma_{X'} = \Sigma'$ and $\mathbf{A}_{X'}$, which can be identified with the subset of colors $D \in \cup_{\alpha\in S\cap\Sigma'}\Delta_X(\alpha)$; one has $c_{X'}(D,\sigma) = c_X(D,\sigma)$ for all $\sigma \in \Sigma'$.

In particular, any spherical root of X is the spherical root of a wonderful G-variety of rank 1. The wonderful G-varieties of rank 1 are well known, for all G. The finite set of spherical roots of wonderful G-varieties of rank 1 is denoted by $\Sigma(G)$. Any spherical root σ is a linear combination of simple roots with non-negative integer coefficients. Spherical roots are usually labeled depending on these coefficients, and we will follow the notation of [2].

We define $\mathrm{supp}\,\sigma$ the set of simple roots whose coefficient in the expression of σ is non-zero, and if $\Sigma' \subseteq \Sigma_X$ then $\mathrm{supp}\,\Sigma'$ is the union of $\mathrm{supp}\,\sigma$ for all $\sigma \in \Sigma'$.

The spherical system of X is determined by the spherical systems of all the localizations $X_{\Sigma'}$ of rank 2 (actually, it is enough to restrict to the localizations of rank 1 and those of rank 2 with a simple spherical root). Furthermore, the wonderful G-varieties of rank 2 are known, for all G.

2.2. **Spherical systems.**

Definition 2.2.1. Let $(S^p, \Sigma, \mathbf{A})$ be a triple such that $S^p \subset S$, $\Sigma \subset \Sigma(G)$ and $\mathbf{A}$ a finite set endowed with a pairing $c\colon \mathbb{Z}\mathbf{A} \times \mathbb{Z}\Sigma_X \to \mathbb{Z}$. For every $\alpha \in \Sigma \cap S$, let $\mathbf{A}(\alpha)$

denote the set $\{D \in \mathbf{A} \colon c(D,\alpha)=1\}$. Such a triple is called a *spherical system* for G if:

- (A1) for every $D \in \mathbf{A}$ we have $c(D,-) \leq 1$, and if $c(D,\sigma)=1$ for some $\sigma \in \Sigma$ then $\sigma \in S \cap \Sigma$;
- (A2) for every $\alpha \in \Sigma \cap S$, $\mathbf{A}(\alpha)$ contains two elements and denoting with D^+_α and D^-_α these elements, it holds $c(D^+_\alpha,-)+c(D^-_\alpha,-)=\langle \alpha^\vee,-\rangle$;
- (A3) the set $\mathbf{A}$ is the union of $\mathbf{A}(\alpha)$ for all $\alpha \in \Sigma \cap S$;
- (Σ1) if $2\alpha \in \Sigma \cap 2S$ then $\frac{1}{2}\langle \alpha^\vee,\sigma\rangle$ is a non-positive integer for every $\sigma \in \Sigma \setminus \{2\alpha\}$;
- (Σ2) if $\alpha,\beta \in S$ are orthogonal and $\alpha+\beta \in \Sigma$ then $\langle \alpha^\vee,\sigma\rangle = \langle \beta^\vee,\sigma\rangle$ for every $\sigma \in \Sigma$;
- (S) for every $\sigma \in \Sigma$, there exists a wonderful G-variety X of rank 1 with $S^p_X = S^p$ and $\Sigma_X = \{\sigma\}$.[1]

The cardinality of Σ is the *rank* of the spherical system.

2.3. **The set of all colors.** Given an abstract spherical system $\mathscr{S} = (S^p,\Sigma,\mathbf{A})$ for a given adjoint group G, it is possible to define the set $\Delta \supseteq \mathbf{A}$ of its colors (and a corresponding extension of c), which plays the role of Δ_X. Following [14] and its notation, we identify each element in $\Delta \setminus \mathbf{A}$ with the simple roots it is moved by. This gives a disjoint union:

$$\Delta = \mathbf{A} \cup \Delta^{2a} \cup \Delta^b$$

Colors in the set Δ^{2a} correspond to simple roots α such that 2α is a spherical root. For such a color D, we have $c(D,-)=\frac{1}{2}\langle \alpha^\vee,-\rangle$. Colors in Δ^b correspond to:

$$\left(S \setminus \left(\Sigma \cup \frac{1}{2}\Sigma \cup S^p\right)\right) \Big/ \sim$$

where $\alpha \sim \beta$ if $\alpha=\beta$, or if $\alpha \perp \beta$ and $\alpha+\beta \in \Sigma$. For such a color D, we have $c(D,-)=\langle \alpha^\vee,-\rangle$, for α any representative of the $\sim$-equivalence class associated to D.

The abstract definition of $\Delta(\alpha)$ for $\alpha \in S$, called again the set of colors *moved* by α, goes as follows:

- if $\alpha \in S \cap \Sigma$, then $\Delta(\alpha) = \{D \in \mathbf{A} \mid c(D,\alpha)=1\}$,
- otherwise $\Delta(\alpha)$ contains at most one element, which is the color corresponding to α following the definitions of Δ^{2a} and Δ^b.

Finally, we say that a color D is *free* if there exists at most one spherical root σ such that $c(D,\sigma)>0$. If $D \in \mathbf{A}$, it is free if and only if it is moved by only one simple root.

2.4. **Quotients.** Certain classes of morphisms between wonderful varieties can be represented by subsets of colors of the domain. Let $f\colon X \to Y$ be a surjective G-equivariant morphism between two wonderful G-varieties, and define $\Delta_f = \{D \in \Delta_X \mid f(D)=Y\}$. Let now Δ' be a subset of Δ_X; we say that Δ' is *distinguished* if there exists a linear combination D of elements of Δ' with positive coefficients, such that $c_X(D,\sigma) \geq 0$ for all spherical roots σ.

We say that Δ' is *(*)-distinguished*, if it is distinguished and the monoid $\{\sigma \in \mathbb{N}\Sigma_X \mid c_X(D,\sigma)=0\ \forall D \in \Delta'\}$ is free; we denote in this case its indecomposable elements by Σ_X/Δ'.

---

[1] This axiom can actually be expressed in purely combinatorial terms.

Proposition 2.4.1 ([14])**.** *Let X be a wonderful G-variety. The map $f \mapsto \Delta_f$ induces a bijection between the set of (*)-distinguished subsets of Δ_X and the set of equivalence classes of couples (f, Y) where $f\colon X \to Y$ is a G-equivariant, surjective map with connected fibers, and where the equivalence relation is induced naturally by G-equivariant isomorphisms $Y \to Y'$.*

Recalling that an inclusion $K \supseteq H$ is *co-connected* if and only if the quotient K/H is connected, we rephrase Proposition 2.4.1 as follows.

Proposition 2.4.2 ([14])**.** *Let X be a wonderful G-variety with generic stabilizer H. For any subgroup K of G containing H, we define $\Delta_K = \Delta_f$ where $f : G/H \to G/K$ is the natural projection. Then $K \mapsto \Delta_K$ is a bijection between the co-connected inclusions $K \supseteq H$ such that $[\mathrm{N}_G K : K]$ is finite and the set of distinguished subsets of Δ_X. The homogeneous space G/K admits a wonderful compactification if and only if Δ_K is (*)-distinguished.*

A (*)-distinguished subset $\Delta' \subseteq \Delta_X$ is

- *smooth* if $\Sigma_X/\Delta' \subseteq \Sigma_X$;
- *homogeneous* (or *parabolic*) if $\Sigma_X/\Delta' = \emptyset$.

Proposition 2.4.3 ([14])**.** *A G-equivariant, surjective map with connected fibers between wonderful varieties $f : X \to Y$ is smooth if and only if Δ_f is smooth. Moreover, Y is homogeneous if and only if Δ_f is homogeneous.*

Obviously, all the above definitions can also be given for an abstract spherical system.

Definition 2.4.1. Let $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$ be a spherical system with set of colors Δ, and let $\Delta' \subseteq \Delta$ be a distinguished subset of colors. We define:

- $S^p/\Delta' = \{\alpha \in S \mid \Delta(\alpha) \subseteq \Delta'\}$;
- Σ/Δ' = the indecomposable elements (or equivalently the minimal generators) of the monoid $\{\sigma \in \mathbb{N}\Sigma \mid c_X(D, \sigma) = 0\ \forall D \in \Delta'\}$;
- $\mathbf{A}/\Delta' = \bigcup_{\alpha \in S \cap \Sigma/\Delta'} \mathbf{A}(\alpha)$.

We say that the quotient $\mathscr{S}/\Delta'$ *exists* if and only if the monoid $\{\sigma \in \mathbb{N}\Sigma \mid c_X(D, \sigma) = 0\ \forall D \in \Delta'\}$ is free and $\mathscr{S}/\Delta' = (S^p/\Delta', \Sigma/\Delta', \mathbf{A}/\Delta')$ is a spherical system. In this case, the bilinear pairing c for $\mathscr{S}/\Delta'$ is induced from that of $\mathscr{S}$ in a natural way.

It is relevant to point out that all distinguished sets of colors are (*)-distinguished, and the quotient $\mathscr{S}/\Delta'$ always exists, but the only proof we know relies on a case-by-case checking ([2]). For this reason we have chosen not to assume it in this paper.

If $\mathscr{S}$ is geometrically realizable and Δ' is (*)-distinguished, then the quotient $\mathscr{S}/\Delta'$ is the spherical system of the wonderful variety Y in the notation of Proposition 2.4.1 (where $\Delta' = \Delta_f$).

2.5. **Localizations.** Let X be a wonderful variety and S' a subset of S. Consider a Levi subgroup $L_{S'}$ of the parabolic subgroup $P_{S'}$; the variety X contains a well defined wonderful $L_{S'}$-subvariety $X_{S'}$, called the *localization* of X in S' (see [14]). Its spherical system satisfies: $S^p_{X_{S'}} = S^p_X \cap S'$; the set $\Sigma_{X_{S'}}$ is the set of spherical roots of X whose support is contained in S'; the set $\mathbf{A}_{X_{S'}}$ is the union of $\Delta_X(\alpha)$ for all $\alpha \in S' \cap \Sigma_X$. The localization $\mathscr{S}_{S'}$ of an abstract spherical system $\mathscr{S}$ in S' is defined analogously.

Recall that we have also defined the localization $X_{\Sigma'}$ of X on a subset of spherical roots[2] $\Sigma' \subseteq \Sigma_X$. The same can be done for an abstract spherical system $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$ and any $\Sigma' \subseteq \Sigma$: we define $\mathscr{S}_{\Sigma'} = (S^p, \Sigma', \mathbf{A}_{\Sigma'})$ where:

$$\mathbf{A}_{\Sigma'} = \bigcup_{\alpha \in S \cap \Sigma'} \Delta(\alpha),$$

and the bilinear pairing on $\mathscr{S}_{\Sigma'}$ is the one of $\mathscr{S}$ restricted to $\mathbb{Z}\Sigma'$.

Given a subset Δ' of the colors Δ of $\mathscr{S}$, we can also define the *restriction* of Δ' to a localization $\mathscr{S}_{S'}$ and to a localization $\mathscr{S}_{\Sigma'}$. In the first case, we define:

$$\Delta'|_{S'} = \Delta' \cap \left(\bigcup_{\alpha \in S'} \Delta(\alpha) \right).$$

It is immediate to show that if $\mathscr{S} = \mathscr{S}_X$ is geometrically realizable, then:

$$\Delta'|_{S'} = \{ D \cap X_{S'} \mid D \in \Delta' \} \subseteq \Delta_{X_{S'}}.$$

In the second case, a more careful analysis is needed, since the full set of colors of a localization $\mathscr{S}_{\Sigma'}$ is not related to the colors of $\mathscr{S}$ as easily as its subset $\mathbf{A}_{\Sigma'}$.

For notational convenience, let us denote by Δ^b_1 the set of $\sim$-equivalence classes containing only one element, and Δ^b_2 those containing two elements. We also write $\Delta_{\Sigma'}$ for the whole set of colors of the spherical system $\mathscr{S}_{\Sigma'}$, and correspondingly $\Delta_{\Sigma'}(\alpha)$ denotes the colors moved by $\alpha \in S$ in the spherical system $\mathscr{S}_{\Sigma'}$.

Then we define the restriction of Δ' to $\mathscr{S}_{\Sigma'}$ as:

$$\Delta'|_{\Sigma'} = \left(\Delta'_{\Sigma',1}\right) \cup \left(\Delta'_{\Sigma',2}\right) \cup \left(\Delta'_{\Sigma',3}\right) \cup \left(\Delta'_{\Sigma',4}\right),$$

where:

$$\Delta'_{\Sigma',1} = \left(\bigcup_{\alpha \in \Sigma' \cap S} \Delta(\alpha) \cap \Delta' \right) \cup \left(\bigcup_{\substack{\alpha \in (\Sigma \setminus \Sigma') \cap S \\ \text{with } \Delta(\alpha) \subseteq \Delta'}} \Delta_{\Sigma'}(\alpha) \right),$$

$$\Delta'_{\Sigma',2} = \bigcup_{\substack{\alpha \in \frac{1}{2}\Sigma \cap S \\ \text{with } \Delta(\alpha) \subseteq \Delta'}} \Delta_{\Sigma'}(\alpha),$$

$$\Delta'_{\Sigma',3} = \bigcup_{\substack{\{\alpha\} \in \Delta^b_1 \\ \text{with } \Delta(\alpha) \subseteq \Delta'}} \Delta_{\Sigma'}(\alpha),$$

$$\Delta'_{\Sigma',4} = \bigcup_{\substack{\{\alpha,\beta\} \in \Delta^b_2 \\ \text{with } \Delta(\alpha) = \Delta(\beta) \subseteq \Delta'}} \left(\Delta_{\Sigma'}(\alpha) \cup \Delta_{\Sigma'}(\beta) \right).$$

This definition is given with the idea of taking a G-equivariant map $f \colon X \to Y$ with connected fibers and study its restriction $f'|_X \colon X' \to f(X')$ to a wonderful subvariety $X' \subset X$. The following proposition and corollary explain this relationship.

---

[2] In cases where $\Sigma' = S' \subseteq \Sigma \cap S$ our notation may be ambiguous: the context will then indicate whether we are localizing to a subset of simple roots or of spherical roots.

Proposition 2.5.1. *Let* Δ' *be a distinguished set of colors of* $\mathscr{S}$ *such that* $\mathscr{S}/\Delta'$ *exists. Then the restriction* $\Delta'|_{\Sigma'}$ *is such that the quotient* $(\mathscr{S}_{\Sigma'})/(\Delta'|_{\Sigma'})$ *exists, and we have:*

$$(\mathscr{S}/\Delta')_{\Sigma''} = (\mathscr{S}_{\Sigma'})/(\Delta'|_{\Sigma'}),$$

where Σ'' *consists of the elements of* Σ/Δ' *that are linear combination of elements of* Σ'.

Proof. Let us fix $E = \sum a_D\, D$, for $D \in \Delta'$, with positive coefficients such that $c(E, \sigma) \geq 0$ for all $\sigma \in \Sigma$, and define $F = \sum b_D\, D$, for $D \in \Delta'|_{\Sigma'}$, as follows (S_D is the set of simple roots moving $D \in \mathbf{A}$):

$$b_D = \begin{cases} \frac{|S_D \cap \Sigma'|}{|S_D|} a_D & \text{if } D \in \Delta_{\Sigma'}(\alpha) \text{ and } \alpha \in \Sigma' \cap S \\ \frac{1}{|S_{D^+_\alpha}|} a_{D^+_\alpha} + \frac{1}{|S_{D^-_\alpha}|} a_{D^-_\alpha} & \text{if } D \in \Delta_{\Sigma'}(\alpha) \text{ and } \alpha \in (\Sigma \setminus \Sigma') \cap S \\ a_D & \text{if } D \in \Delta_{\Sigma'}(\alpha) \text{ and } \alpha \in \frac{1}{2}\Sigma' \cap S \\ \frac{1}{2} a_D & \text{if } D \in \Delta_{\Sigma'}(\alpha) \text{ and } \alpha \in \frac{1}{2}(\Sigma \setminus \Sigma') \cap S \\ a_D & \text{if } D \in \Delta_{\Sigma'}(\alpha) \text{ and } \alpha \in \Delta^b_1 \\ a_D & \text{if } D \in \Delta_{\Sigma'}(\alpha) \cap \Delta_{\Sigma'}(\beta) \text{ and } \{\alpha, \beta\} \in \Delta^b_2 \\ \frac{1}{2} a_D & \text{if } D \in \Delta_{\Sigma'}(\alpha) \setminus \Delta_{\Sigma'}(\beta) \text{ and } \{\alpha, \beta\} \in \Delta^b_2 \end{cases}$$

For all $\sigma \in \Sigma'$ one has $c(F, \sigma) = c(E, \sigma) \geq 0$, therefore $\Delta'|_{\Sigma'}$ is distinguished in $\mathscr{S}_{\Sigma'}$.

The equality, without assuming that the triple $(\mathscr{S}_{\Sigma'})/(\Delta'|_{\Sigma'})$ is a spherical system, is easily checked. Finally, the existence of the quotient $(\mathscr{S}_{\Sigma'})/(\Delta'|_{\Sigma'})$ follows from the existence of $\mathscr{S}/\Delta'$. □

Corollary 2.5.1. *Let* $f\colon X \to Y$ *be a* G*-equivariant morphism with connected fibers between wonderful varieties, and let* $X' \subseteq X$ *be a* G*-stable subvariety. Then* $Y' = f(X')$ *is a* G*-stable subvariety of* Y*, and:*

$$\mathscr{S}_{Y'} = \mathscr{S}_{X'}/(\Delta_f|_{\Sigma_{X'}}).$$

Definition 2.5.1. A spherical system $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$ is *cuspidal* if and only if $\operatorname{supp} \Sigma = S$.

Lemma 2.5.1 ([14])**.** *Let* $\mathscr{S}$ *be a geometrically realizable spherical system with generic stabilizer* H*, and suppose that* H *does not contain any simple factor of* G*. Then* $\mathscr{S}$ *is not cuspidal if and only if there exists a proper parabolic subgroup* P *of* G *such that* $P^r \subseteq H \subseteq P$.

If $\mathscr{S}$ is not cuspidal, then its wonderful variety is obtained from its localization to $(S^p \cup \operatorname{supp} \Sigma) \subset S$ by a procedure called *parabolic induction*: see [14] for details.

3. From the spherical system to the wonderful subgroup

3.1. The Levi part.

Let $\mathscr{S}$ be a spherical system. The problem of determining a Levi part of a generic stabilizer H of $\mathscr{S}$ (or a candidate, if geometric realizability of $\mathscr{S}$ is not assured) is related to a special class of spherical subgroups. These subgroups have the form $K = L_K P^u$, where P is a parabolic subgroup containing K, and L_K is a very reductive spherical subgroup of a Levi part L_P of P.

If $\mathscr{S}$ is geometrically realizable, then it admits a subset $\Delta' \subseteq \Delta$ corresponding to a subgroup K of the above form. More precisely, Δ' can be found in such a way that L_K and a Levi component of H differ only by their connected center.

The combinatorial version of these notions is the following.

Definition 3.1.1. Let $\mathscr{S}$ be a spherical system with colors Δ. A couple of subsets (Δ',Δ'') with $\Delta'\subseteq\Delta''\subseteq\Delta$ is of type $(\overline{\mathscr{L}},\overline{\mathscr{R}})$ if and only if Δ' is distinguished, Δ'' is minimal parabolic, and they satisfy the following conditions:

1. no simple root in the support of Σ/Δ' moves a color in $\Delta\setminus\Delta''$;
2. there exists a linear combination of elements in Σ/Δ' with positive coefficients, such that it takes non-negative values on all colors in $\Delta''\setminus\Delta'$;
3. Δ' is minimal with these properties.

We also say that a subset of colors $\Delta'\subseteq\Delta$ is of type $(\overline{\mathscr{L}})$ if there exists some Δ'' such that the couple (Δ',Δ'') is of type $(\overline{\mathscr{L}},\overline{\mathscr{R}})$.

Proposition 3.1.1. *Suppose that the spherical system $\mathscr{S}$ is geometrically realizable, with generic stabilizer H and set of colors Δ. Then subsets $\Delta'\subseteq\Delta$ of type $(\overline{\mathscr{L}})$ are in bijection with co-connected $K\supseteq H$ with the following properties:*

1. *there exists a parabolic subgroup P minimal containing H, such that $P\supseteq K\supseteq P^u$;*
2. *there exist Levi components $L_P\supseteq L_K\supseteq L$ of resp. P, K and H, such that:*
   (a) *L_K and L are very reductive subgroups of L_P, so in particular $K=L_KP^u$,*
   (b) *$Z(L)^\circ\subseteq Z(L_K)^\circ$ and:*

$$\frac{L}{Z(L)^\circ}=\frac{L_K}{Z(L_K)^\circ}.$$

Proof. If we start with a co-connected inclusion $P\supseteq K\supseteq H$ with the above hypotheses, it corresponds to a minimal parabolic subset Δ'' and a subset $\Delta'\subseteq\Delta''$ of type $(\overline{\mathscr{L}})$. Indeed, condition (1) of Definition 3.1.1 follows from the fact that G/K is the parabolic induction by means of P of the homogeneous space L_P/L_K, and condition (2) follows from the fact that L_K is reductive in L_P.

Let (Δ',Δ'') be of type $(\overline{\mathscr{L}},\overline{\mathscr{R}})$. Let $K\supseteq H$ and $P\supseteq K\supseteq H$ be the corresponding co-connected inclusions. From Definition 3.1.1 it follows that $\mathscr{S}/\Delta'$ is the parabolic induction of a reductive spherical system, by means of the parabolic subgroup P, and claim (1) is proved.

The minimality of P containing H implies that we can choose Levi components $L_P\supseteq L_K\supseteq L$ with L very reductive in L_P. Hence L_K is very reductive in L_P too: this implies $Z(L)^\circ\subseteq Z(L_K)^\circ$.

If:

$$\left(\frac{L}{Z(L)^\circ}\right)^\circ\neq\left(\frac{L_K}{Z(L_K)^\circ}\right)^\circ$$

then the subgroup $\widetilde{K}=LP^r$ is strictly contained in K. The subgroup $P\supseteq\widetilde{K}'\supseteq H$ satisfies the conditions of the statement. The corresponding subset of colors $\widetilde{\Delta'}\subsetneq\Delta'$ satisfies conditions (1) and (2), by the first part of the proof, thus contradicting the minimality of Δ'.

We conclude that:

$$\left(\frac{L}{Z(L)^\circ}\right)^\circ=\left(\frac{L_K}{Z(L_K)^\circ}\right)^\circ,$$

and the equality:

$$\frac{L}{Z(L)^\circ}=\frac{L_K}{Z(L_K)^\circ}$$

then follows from the fact that $K\supseteq H$ is co-connected. □

The above proposition gives a procedure to determine a Levi part of H, up to its connected center, starting from $\mathscr{S}$. Namely, let Δ' be a subset of colors of type $(\overline{\mathscr{L}})$. The quotient $\mathscr{S}/\Delta'$ is the parabolic induction of a reductive spherical system $\mathscr{T}$. Once $\mathscr{T}$ is computed, its generic stabilizer L_K is obtained from the classification of reductive wonderful subgroups and their spherical systems.

3.2. **The connected center.**

Let $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$ be a geometrically realizable spherical system with generic stabilizer H and set of colors Δ.

Conjugating H if necessary, let us choose a parabolic subgroup Q_- containing B_- and H, such that Q_- is minimal containing H. The subgroup Q_- is associated to a distinguished minimal parabolic subset of colors Δ_{Q_-}.

From minimality, it follows that the projection of H in Q_-/Q_-^u is not contained in any parabolic subgroup of Q_-/Q_-^u. This projection is therefore reductive, or equivalently: $Q_-^u \supseteq H^u$.

Recall that L_{Q_-} denotes the standard Levi subgroup of Q_-, and let us choose a Levi part $L \subseteq L_{Q_-}$. The considerations above tell us that L is very reductive inside L_{Q_-}, and that $Z(L)^\circ$ is a subtorus of $Z(L_{Q_-})^\circ \subseteq T$.

The dimension of $Z(L)^\circ$ is immediately given by the well known formula:

$$\dim Z(L)^\circ = \operatorname{card}\Delta - \operatorname{card}\Sigma.$$

This integer is also called the *defect* of $\mathscr{S}$, and denoted by $\mathrm{d}(\mathscr{S})$.

Definition 3.2.1. We define the following subsets of $\mathbb{Z}\Sigma$:

$$\rho(\Delta_{Q_-})^\perp = \{\gamma \in \mathbb{Z}\Sigma \mid c(D,\gamma) = 0 \ \forall D \in \Delta_{Q_-}\}$$

$$\rho(\Delta_{Q_-})^\vee = \{\gamma \in \mathbb{Z}\Sigma \mid c(D,\gamma) \geq 0 \ \forall D \in \Delta_{Q_-}\}.$$

Obviously, $\rho(\Delta_{Q_-})^\perp$ is a sublattice of $\mathbb{Z}\Sigma$.

For each color D in $\Delta \setminus \Delta_{Q_-}$ define α_D as the unique simple root moving D (this root is unique because Δ_{Q_-} is homogeneous). Define also λ_D to be the fundamental dominant weight associated to α_D, and $\lambda_D^* = -w_0(\lambda_D)$, where w_0 is the longest element in the Weyl group.

Finally, for all $\gamma \in \rho(\Delta_{Q_-})^\perp$, define $\mu(\gamma)$ to be the weight:

$$\sum_{D\in\Delta\setminus\Delta_{Q_-}} c(D,\gamma)\cdot\lambda_D^*$$

restricted to the subgroup $Z(L_{Q_-})^\circ$ of T.

Lemma 3.2.1. *The map $\gamma \mapsto \mu(\gamma)$ is an isomorphism:*

$$\mu\colon \rho(\Delta_{Q_-})^\perp \to \mathcal{X}\left(Z(L_{Q_-})^\circ\right)^{Z(L)\cap Z(L_{Q_-})^\circ}$$

Proof. Let $\gamma \in \rho(\Delta_{Q_-})^\perp$ and consider a B-proper rational function f_γ on G/H having B-weight γ. Call F_γ the pull-back of f_γ on G along the projection $G \to G/H$.

Since f_γ has neither zeros nor poles on colors of Δ_{Q_-}, F_γ has neither zeros nor poles on the pull-backs on G of these colors. Its zeros or poles must then lie only on the pullbacks on G of colors of G/Q_- along the projection $G \to G/Q_-$.

This means that F_γ is Q_--proper under right translation, and it is immediate to see that the Q_--weight of F_γ is just $\mu(\gamma)$. But F_γ is also of course H-stable under right translation, and this implies that $\mu(\gamma)$ is constantly 1 on $Z(L)\cap Z(L_{Q_-})^\circ$.

Viceversa, suppose to have any weight χ of $Z(L_{Q_-})^\circ$ which is constantly 1 on $Z(L)\cap Z(L_{Q_-})^\circ$: considering χ as a Q_--weight, we can write it as a linear combination with integer coefficients of weights of the form λ^*_D for D ranging through the colors of G/Q_-:

$$\chi = \sum_{D\in\Delta\setminus\Delta_{Q_-}} a_D\cdot\lambda^*_D.$$

So χ is the Q_--weight (under right translation) of some rational function on G, call it F, which is a B-eigenvector under left translation and a Q_--eigenvector under right translation, whose zeros and poles lie on the pullbacks on G of colors of G/Q_- along the projection $G\to G/Q_-$.

The function F under right translation is Q_-^u-stable and (L_{Q_-},L_{Q_-})-stable; it is a $Z(L_{Q_-})^\circ$-eigenvector but $Z(L)^\circ$-stable, since $Z(L)^\circ\subseteq Z(L)\cap Z(L_{Q_-})^\circ$. Thus F is H-stable, and it descends to a rational function f on G/H. Having neither zeros nor poles along colors in Δ_{Q_-} by construction, the function f is equal to f_γ for some $\gamma\in\rho(\Delta_{Q_-})^\perp$.

In addition, we notice that the weights λ^*_D of $Z(L_{Q_-})^\circ$ are independent (over $\mathbb{Q}$). Indeed, the colors in $\Delta\setminus\Delta_{Q_-}$ correspond each to a different simple root, that we called α_D, and these simple roots are exactly those which are not associated to Q_-.

Finally, we can reapply the first part of the proof to $f=f_\gamma$: since the Q_--weight (by right translation) of F is χ, we have $a_D=c(D,\gamma)$ for all $D\in\Delta\setminus\Delta_{Q_-}$. As a consequence, $\mu(\gamma)=\chi$ and the proof is complete. ☐

Corollary 3.2.1. *We have the following equality between subgroups of* $Z(L_{Q_-})^\circ$*:*

$$Z(L)^\circ = \left(\bigcap_{\gamma\in\rho(\Delta_{Q_-})^\perp}\ker\mu(\gamma)\right)^\circ.$$

Proof. It directly follows from Lemma 3.2.1. ☐

If we start with an abstract spherical system $\mathscr{S}$ and a minimal parabolic subset of colors, we can repeat this procedure just using $Z(L_{Q_-})^\circ$ and $\mathscr{S}$. A candidate connected center $Z(L)^\circ$ is therefore always defined.

Corollary 3.2.2. *The codimension of* $Z(L)^\circ$ *inside* $Z(L_{Q_-})^\circ$ *equals the codimension of the convex cone* $\mathbb{Q}_{\geq 0}\rho(\Delta_{Q_-})$ *inside* $\mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}\Sigma,\mathbb{Q})$.

Proof. From Lemma 3.2.1 and Corollary 3.2.1 we have that the rank of the lattice $\rho(\Delta_{Q_-})^\perp$ is equal to the codimension of $Z(L)^\circ$ inside $Z(L_{Q_-})^\circ$.

It remains to notice that $\rho(\Delta_{Q_-})^\perp$ is the linear part of the convex cone $\rho(\Delta_{Q_-})^\vee$, and that the latter is the dual of the convex cone $\mathbb{Q}_{\geq 0}\rho(\Delta_{Q_-})$. ☐

3.3. **The unipotent radical.** We maintain the assumptions of the previous section. We know that $\mathrm{Lie}\,Q_-^u/\mathrm{Lie}\,H^u$ is a spherical L-module. The following proposition follows from the results of [12].

Proposition 3.3.1 ([12])**.** *The spherical system* $\mathscr{S}$ *uniquely determines the structure of* $\mathrm{Lie}\,Q_-^u/\mathrm{Lie}\,H^u$ *as an* L*-module.*

However, there is no known direct technique to compute of $\mathrm{Lie}\,Q_-^u/\mathrm{Lie}\,H^u$ from $\mathscr{S}$ unless one uses case-by-case considerations, for example based on the list of affine smooth spherical varieties of [10].

On the other hand, the spherical system provides immediate information on the fiber Q_-/H of the map $G/H \to G/Q_-$. Our assumptions imply that Q_-/H is an affine smooth variety, spherical under the action of L_{Q_-} (see [4]), and L_{Q_-}-equivariantly isomorphic to the homogeneous vector bundle $L_{Q_-} \times_L \mathrm{Lie}\, Q_-^u / \mathrm{Lie}\, H^u$.

Definition 3.3.1. Recall that $B \cap L_{Q_-}$ is a Borel subgroup of L_{Q_-}. We denote by $\Lambda^+(Q_-/H)$ the monoid of $(B \cap L_{Q_-})$-eigenvalues of elements in $\mathbb{C}[Q_-/H]^{(B\cap L_{Q_-})}$.

Lemma 3.3.1. *We have* $\Lambda^+(Q_-/H) = \rho(\Delta_{Q_-})^\vee$.

Proof. Let $f \in \mathbb{C}(G/H)^{(B)}$, and consider its restriction $f|_F$ on the fiber $F = Q_-/H \subseteq G/H$. The restriction is a $(B \cap L_{Q_-})$-eigenvector, and it is non-zero because Q_- is B-spherical, therefore no color of G/H contains F.

We obtain in this way a bijection between $\mathbb{C}(G/H)^{(B)}$ and $\mathbb{C}(F)^{(B\cap L_{Q_-})}$. This follows from [6, Proposition I.2]; let us provide here a direct proof. Injectivity is obvious, since T is a maximal torus of $(B \cap L_{Q_-})$, therefore the $(B \cap L_{Q_-})$-eigenvalue of $f|_F$ is equal to the B-eigenvalue of f. Let now $f_0 \in \mathbb{C}(Q_-/H)^{(B\cap L_{Q_-})}$, and let us extend it to a function $f \in \mathbb{C}(G/H)^{(B)}$ with the following definition:

$$f(uqH) = f_0(qH)$$

for $u \in B^u$ and $q \in Q_-$. It is easy to show that f is a well defined rational function on the open set $B^u Q_-/H$ of G/H, and that it is a B-eigenvector.

Now, functions in $\mathbb{C}[F]^{(B\cap L_{Q_-})}$ are exactly the restrictions of functions on G/H having no pole on colors in Δ_{Q_-}, and the lemma follows. □

A further step, going beyond the scope of the present work, would be to study all possible spherical quotients of $\mathrm{Lie}\, Q_-^u$ under the action of spherical very reductive subgroups of L_{Q_-}. Taking into account the results of §5, it would be enough to assume that the very reductive subgroup contains $Z(L_{Q_-})^\circ$: this would provide a strategy to guess H^u inside Q_-^u based only on $\mathscr{S}$.

4. Wonderful fiber products

4.1. Let $X_1, X_2, X_{1,2}$ be wonderful G-varieties and $\varphi_1 \colon X_1 \to X_{1,2}$, $\varphi_2 \colon X_2 \to X_{1,2}$ surjective equivariant maps with connected fibers. Then the fiber product $X = X_1 \times_{X_{1,2}} X_2$ is a (not necessarily spherical) G-variety, with $\psi_1 \colon X \to X_1$, $\psi_2 \colon X \to X_2$ surjective equivariant maps with connected fibers such that $\varphi_1 \circ \psi_1 = \varphi_2 \circ \psi_2$.

Definition 4.1.1. Let $X_1, X_2, X_{1,2}$ be wonderful G-varieties with maps as above. Let $Z_1, Z_2, Z_{1,2}$ be the corresponding closed G-orbits, with the corresponding restricted maps. The G-variety $X = X_1 \times_{X_{1,2}} X_2$ is called wonderful fiber product if it is wonderful with closed G-orbit $Z \cong Z_1 \times_{Z_{1,2}} Z_2$.

On the combinatorial level we give the following.

Definition 4.1.2. Let $\mathscr{S}$ be a spherical system. Two (*)-distinguished subsets of colors Δ_1, Δ_2 are said to decompose $\mathscr{S}$ if:

- $(S^p/\Delta_1 \setminus S^p) \perp (S^p/\Delta_2 \setminus S^p)$,
- $\Sigma \subset (\Sigma/\Delta_1 \cup \Sigma/\Delta_2)$.

If Δ_1, Δ_2 decompose $\mathscr{S}$ then there is no $\sigma \in \Sigma$, $D_1 \in \Delta_1$, $D_2 \in \Delta_2$ such that both $c(D_1, \sigma)$ and $c(D_2, \sigma)$ are non-zero. Therefore, $\mathrm{card}(\Sigma) + \mathrm{card}(\Sigma/\Delta_1 \cup \Delta_2) = \mathrm{card}(\Sigma/\Delta_1) + \mathrm{card}(\Sigma/\Delta_2)$. Moreover, $P_{S^p} = P_{S^p/\Delta_1} \cap P_{S^p/\Delta_2}$ and $P_{S^p/(\Delta_1 \cup \Delta_2)} = P_{S^p/\Delta_1} P_{S^p/\Delta_2} = P_{S^p/\Delta_2} P_{S^p/\Delta_1}$.

Theorem 4.1.1. *Let X be a wonderful G-variety with spherical system $\mathscr{S}$. Then X is isomorphic to a wonderful fiber product $X_1 \times_{X_{1,2}} X_2$ (with notation as above) if and only if the corresponding (*)-distinguished subsets $\Delta_{\psi_1}, \Delta_{\psi_2}$ decompose $\mathscr{S}$.*

Proof. Recall the local structure of a wonderful G-variety X: $X_B = X \setminus \cup_{D \in \Delta} D$ is an open affine B-subvariety P_{S^p}-isomorphic to $P^u_{S^p} \times M$, where M is a smooth affine T-subvariety isomorphic to $\mathbb{C}^\Sigma$. A surjective G-equivariant map with connected fibers between wonderful G-varieties $X \to X'$ restricts to a B-equivariant map between affine B-varieties $X_B \to X'_B$ and, moreover, to a T-equivariant map between affine T-varieties $\mathbb{C}^\Sigma \to \mathbb{C}^{\Sigma/\Delta'}$ corresponding to the inclusion $\mathbb{N}\,\Sigma/\Delta' \subset \mathbb{N}\,\Sigma$.

Therefore, if X is wonderful fiber product then one has

$$\mathbb{C}^\Sigma = \mathbb{C}^{\Sigma/\Delta_1} \times_{\mathbb{C}^{\Sigma/\Delta_1 \cup \Delta_2}} \mathbb{C}^{\Sigma/\Delta_2} \tag{4.1}$$

which together with the condition on the closed orbits

$$Z \cong Z_1 \times_{Z_{1,2}} Z_2 \tag{4.2}$$

is equivalent to the conditions given in Definition 4.1.2.

Vice versa, once one has (4.1) and (4.2), then consider the fiber product $X' = X_1 \times_{X_{1,2}} X_2$. It contains the open affine B-subvariety $(X_1)_B \times_{(X_{1,2})_B} (X_2)_B$ which has an open B-orbit: then X' is spherical. Furthermore, it is: smooth since $M' = \mathbb{C}^\Sigma$ is; complete; and toroidal since every color of X' maps non-dominantly into X_1 or X_2, which are both toroidal. Thence, X' is wonderful and isomorphic to X. □

From the above proof, it is clear that we have also the following.

Theorem 4.1.2. *Let $\mathscr{S}$ be a spherical system, and let Δ_1, Δ_2 be (*)-distinguished subsets of colors that decompose $\mathscr{S}$. If $\mathscr{S}/\Delta_1$ and $\mathscr{S}/\Delta_2$ are geometrically realizable then $\mathscr{S}$ is geometrically realizable.*

If $X = X_1 \times_{X_{1,2}} X_2$ is wonderful fiber product, choose x in the open orbit and set $x_1 = \psi_1(x)$, $x_2 = \psi_2(x)$, $x_{1,2} = \varphi_1(x_1) = \varphi_2(x_2)$. Let $H, H_1, H_2, H_{1,2}$ be the corresponding stabilizers, then $H = H_1 \cap H_2$ and $H_{1,2} = H_1 H_2 = H_2 H_1$, namely $G/H \cong G/H_1 \times_{G/H_{1,2}} G/H_2$.

5. Quotients of type ($\mathscr{L}$)

5.1. **Combinatorial characterization of quotients of type ($\mathscr{L}$).** Following [4], we give the following:

Definition 5.1.1. Let H be a wonderful subgroup, and $K \supset H$ be a co-connected inclusion. Then $K \supset H$ is *of type ($\mathscr{L}$)* if and only if H^u is strictly contained in K^u, $\mathrm{Lie}\, K^u / \mathrm{Lie}\, H^u$ is a simple H-module and Levi components of H and K (L and L_K, respectively) differ only by their connected centers (actually, $L = \mathrm{N}_{L_K}(H^u)$).

The combinatorial counterpart of Definition 5.1.1 is:

Definition 5.1.2. Let $\mathscr{S}$ be a spherical system with set of colors Δ. A minimal (non-empty) distinguished subset of colors Δ' is of type ($\mathscr{L}$) if and only if it is contained in subsets $\widetilde{\Delta'}$ and Δ'' such that $(\widetilde{\Delta'}, \Delta'')$ is of type $(\overline{\mathscr{L}}, \overline{\mathscr{R}})$.

Proposition 5.1.1. *Suppose that $\mathscr{S}$ is a geometrically realizable spherical system with generic stabilizer H. Then the bijection of Proposition 2.4.2 restricts to a bijection between distinguished subsets $\Delta' \subseteq \Delta$ of type ($\mathscr{L}$) and co-connected inclusions $K \supset H$ of type ($\mathscr{L}$).*

Proof. If we have a co-connected inclusion $K \supset H$ of type ($\mathscr{L}$), the corresponding subset of colors Δ_K is minimal distinguished, so we have only to show that it is also contained in a subset of type ($\overline{\mathscr{L}}$).

Let P be a parabolic subgroup minimal containing H. Since the inclusion $K \supset H$ is of type ($\mathscr{L}$), up to conjugation we can suppose that $P \supseteq K \supset H$, that $P^u \supseteq K^u \supset H^u$, and that we have chosen Levi parts $L_P \supseteq L_K \supseteq L$ in such a way that L_K and L differ only by their connected centers.

Consider the subgroup $K' = LP^r$. We have $P \supseteq K' \supseteq K \supseteq H$, and one checks using Proposition 3.1.1 that $K' \supseteq H$ corresponds to a minimal subset of colors $\Delta_{K'}$ of type ($\overline{\mathscr{L}}$) containing Δ_K.

Viceversa, suppose we have a subset of type ($\overline{\mathscr{L}}$) containing a minimal subset of colors. From geometric realizability we deduce that they are associated to subgroups $K' \supseteq K \supset H$. We can thus call our subsets of colors resp. $\Delta_{K'}$ and Δ_K.

Since $\Delta_{K'}$ is of type ($\overline{\mathscr{L}}$), we know that Levi parts of H and K' differ only by their connected centers. This must happen for K and H too, since we can suppose to have Levi parts $L_{K'} \supseteq L_K \supseteq L$.

It also follows that $K^u \supset H^u$. The minimality of Δ_K assures that H^u contains (K^u, K^u): indeed, the projection of H^u on $K^u/(K^u, K^u)$ cannot be surjective because otherwise we would have $H^u = K^u$. The minimality then implies $H^u \supseteq (K^u, K^u)$.

Hence $\operatorname{Lie} H^u$ is an ideal of $\operatorname{Lie} K^u$ and $\operatorname{Lie} K^u/\operatorname{Lie} H^u$ is abelian: again the minimality of Δ_K implies that it is a simple L-module. Therefore $K \supset H$ is of type ($\mathscr{L}$). □

5.2. The generic stabilizer of a system admitting a quotient of type ($\mathscr{L}$).

Let $\mathscr{S}$ be a spherical system admitting a quotient $\mathscr{S}/\Delta'$ of type ($\mathscr{L}$), one would like to construct the generic stabilizer H of $\mathscr{S}$ starting from the generic stabilizer of $\mathscr{S}/\Delta'$.

The construction we propose is here below. At present, the only proof of the below conjectures that we can imagine would be case-by-case after reducing to the list of primitive spherical systems.

We introduce an additional hypothesis which makes the description easier and excludes only "a few" special quotients of type ($\mathscr{L}$). A color is called *negative* if is ≤ 0 on the set of spherical roots (in this case it is moved by a unique simple root). A negative color is called *interior* (resp. *exterior*) if is moved by a simple root belonging (resp. not belonging) to the support of a spherical root. We assume that Δ' is minimal, there exist exterior negative colors of $\mathscr{S}/\Delta'$ that are interior or not negative as colors of $\mathscr{S}$ and there exist no interior negative colors of $\mathscr{S}/\Delta'$ that are not negative as colors of $\mathscr{S}$. This implies that Δ' is of type ($\mathscr{L}$), see [4].

Let $S_{\Delta'}$ be the subset of simple roots moving exterior negative colors of $\mathscr{S}/\Delta'$ that are interior or not negative as colors of $\mathscr{S}$. Let Q_- be the parabolic subgroup corresponding to $S \setminus S_{\Delta'}$ and let K be the generic stabilizer of $\mathscr{S}/\Delta'$: we can assume $Q^r_- \subset K \subset Q_-$. Set a decomposition $Q_- = Q^r_- M$, we have $K = Q^r_- K_{\Delta'}$ with $K_{\Delta'} \subset M$ (notice that $K_{\Delta'}$ is not necessarily reductive).

Let us consider the spherical systems $\mathscr{T}$ and $\mathscr{T}'$, of M, obtained by localization of $\mathscr{S}$ and $\mathscr{S}/\Delta'$, resp., in the subset of simple roots $S \setminus S_{\Delta'}$. The spherical system $\mathscr{T}'$ equals the quotient spherical system $\mathscr{T}/\Delta'|_{S\setminus S_{\Delta'}}$. Let $H_{\Delta'}$ and $K_{\Delta'}$ be the corresponding generic stabilizers, $H_{\Delta'} \subset K_{\Delta'} \subset M$.

Conjecture 5.2.1. *There exists a simple $K_{\Delta'}$-module W such that $\mathbb{P}(W) \cong \overline{K_{\Delta'}/H_{\Delta'}}$.*

Let W^* be the dual of W. The space $\mathbb{P}(W^*)$ still has an open spherical $K_{\Delta'}$-orbit (notice that a module is spherical if and only if its dual is). Let $H^*_{\Delta'} \subset K_{\Delta'}$ be the generic stabilizer, namely $\mathbb{P}(W^*) \cong \overline{K_{\Delta'}/H^*_{\Delta'}}$, it corresponds to a spherical system $\mathscr{T}^*$ admitting $\mathscr{T}/\Delta'$ as quotient.

Conjecture 5.2.2. *For all $\alpha \in S_{\Delta'}$ there exists a $K_{\Delta'}$-orbit closure in $\mathbb{P}(V(-\alpha)^*)$ such that $\overline{K_{\Delta'}/H^*_{\Delta'}} \cong \mathbb{P}(W^*) \subset \mathbb{P}(V(-\alpha)^*)$.*

This is equivalent to a combinatorial statement on the spherical system $\mathscr{T}^*$, see [4, §2.4].

Then $\mathrm{Lie}\, H^u \cap \mathrm{Lie}\, Q^u_-$ is the co-simple $K_{\Delta'}$-submodule of $\mathrm{Lie}\, Q^u_-$ dual to W^* diagonally included in $V^* = \oplus_{\alpha \in S_{\Delta'}} V(-\alpha)^*$ (notice that $\mathrm{Lie}\, Q^u_- = V \oplus [\mathrm{Lie}\, Q^u_-, \mathrm{Lie}\, Q^u_-]$).

Therefore, $H = (H^u \cap Q^u_-)\, K_{\Delta'}\, \mathrm{N}_{Z(Q_-)}(H^u \cap Q^u_-)$.

Let us give some examples of quotients of type $(\mathscr{L})$.

5.2.1. Let G be $\mathrm{Sp}(6)$. Consider the spherical system $\mathscr{S}$ with $S^p = \emptyset$, $\Sigma = \{\alpha_1+\alpha_2, \alpha_2+\alpha_3\}$ and $\mathbf{A} = \emptyset$. The set of colors is $\Delta = \{D_{\alpha_1}, D_{\alpha_2}, D_{\alpha_3}\}$ with Cartan pairing as follows.

$c(-,-)$	$\alpha_1+\alpha_2$	$\alpha_2+\alpha_3$
D_{α_1}	1	-1
D_{α_2}	1	0
D_{α_3}	-1	1

The subset $\Delta' = \{D_{\alpha_2}\}$ is distinguished of type $(\mathscr{L})$. The corresponding quotient $\mathscr{S}/\Delta'$ has $S^p/\Delta' = \{\alpha_2\}$, $\Sigma/\Delta' = \{\alpha_2+\alpha_3\}$ and $\mathbf{A} = \emptyset$.



Here $S_{\Delta'} = \{\alpha_2\}$, $K_{\Delta'} \cong \mathrm{SL}(2) \times \mathrm{SL}(2)$, $H_{\Delta'}$ is a codimension 1 parabolic subgroup of $K_{\Delta'}$ and W is isomorphic to the standard representation of one simple factor of $K_{\Delta'}$.

5.2.2. Let G be $\mathrm{Sp}(10)$. Consider the spherical system $\mathscr{S}$ with $S^p = \emptyset$, $\Sigma = S$, $\mathbf{A} = \Delta = \{D^+_{\alpha_1} = D^+_{\alpha_3} = D^+_{\alpha_5}, D^+_{\alpha_2}, D^+_{\alpha_4}, D^-_{\alpha_1} = D^-_{\alpha_4}, D^-_{\alpha_2} = D^-_{\alpha_5}, D^-_{\alpha_3}\}$ and Cartan pairing as follows.

$c(-,-)$	α_1	α_2	α_3	α_4	α_5
$D^+_{\alpha_1}$	1	-1	1	-1	1
$D^+_{\alpha_2}$	0	1	0	0	-1
$D^+_{\alpha_4}$	-1	0	0	1	-1
$D^-_{\alpha_1}$	1	0	-1	1	-1
$D^-_{\alpha_2}$	-1	1	-1	0	1
$D^-_{\alpha_3}$	-1	0	1	0	-1

The subset $\Delta' = \Delta \setminus \{D^-_{\alpha_3}\}$ is distinguished of type $(\mathscr{L})$. The corresponding quotient $\mathscr{S}/\Delta'$ has $S^p/\Delta' = S \setminus \{\alpha_3\}$, $\Sigma/\Delta' = \emptyset$ and $\mathbf{A} = \emptyset$.



In this case $S_{\Delta'} = \{\alpha_3\}$, $Q_- = K$ and $K_{\Delta'} = M \cong \mathrm{SL}(3) \times \mathrm{Sp}(4)$. One gets $W = V(-\alpha_3)$ (this would directly follow from the conjecture by dimension reasons). To find $H_{\Delta'}$ one has to reapply the same construction to a quotient of type $(\mathscr{L})$ of $\mathscr{T}$. Let us reset our notation.

Let G be $\mathrm{SL}(3) \times \mathrm{Sp}(4)$. Consider the spherical system $\mathscr{S}$ with $S^p = \emptyset$, $\Sigma = S$, $\mathbf{A} = \Delta = \{D^+_{\alpha'_1} = D^+_{\alpha''_2}, D^+_{\alpha'_2}, D^+_{\alpha''_1}, D^-_{\alpha'_1} = D^-_{\alpha''_1}, D^-_{\alpha'_2} = D^-_{\alpha''_2}\}$ and Cartan pairing as follows.

$c(-,-)$	α'_1	α'_2	α''_1	α''_2
$D^+_{\alpha'_1}$	1	-1	-1	1
$D^+_{\alpha'_2}$	0	1	0	-1
$D^+_{\alpha''_1}$	-1	0	1	-1
$D^-_{\alpha'_1}$	1	0	1	-1
$D^-_{\alpha'_2}$	-1	1	0	1

The subset $\Delta' = \{D^-_{\alpha'_1}, D^-_{\alpha'_2}\}$ is distinguished of type $(\mathscr{L})$. The corresponding quotient $\mathscr{S}/\Delta'$ has $S^p/\Delta' = \emptyset$, $\Sigma/\Delta' = \{\alpha'_1 + \alpha''_2\}$ and $\mathbf{A} = \emptyset$.



Therefore, $S_{\Delta'} = \{\alpha'_2, \alpha''_1\}$, $M \cong \mathrm{SL}(2) \times \mathrm{SL}(2)$ and $K^\circ_{\Delta'} \cong \mathrm{SL}(2)$ included diagonally in M. Here $H_{\Delta'}$ is a Borel subgroup of $K_{\Delta'}$, W is isomorphic to the standard representation of $K_{\Delta'}$ and $W \cong V(-\alpha'_2) \cong V(-\alpha''_1)$, hence W is included diagonally in $V(-\alpha'_1) \oplus V(-\alpha''_2)$.

5.2.3. Let G be $\mathrm{SO}(7)$. Consider the spherical system $\mathscr{S}$ with $S^p = \emptyset$, $\Sigma = \{\alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_3\}$, $\mathbf{A} = \{D^+_{\alpha_3}, D^-_{\alpha_3}\}$ with $c(D^+_{\alpha_3}, \sigma) = c(D^-_{\alpha_3}, \sigma)$ for all $\sigma \in \Sigma$. Its set of colors is $\Delta = \{D_{\alpha_1}, D_{\alpha_2}, D^+_{\alpha_3}, D^-_{\alpha_3}\}$ with Cartan pairing as follows.

$c(-,-)$	$\alpha_1 + \alpha_2$	$\alpha_2 + \alpha_3$	α_3
D_{α_1}	1	-1	0
D_{α_2}	1	1	-1
$D^+_{\alpha_3}$	-1	0	1
$D^-_{\alpha_3}$	-1	0	1

The subset $\Delta' = \{D_{\alpha_2}, D^+_{\alpha_3}\}$ is distinguished of type $(\mathscr{L})$. The corresponding quotient $\mathscr{S}/\Delta'$ has $S^p/\Delta' = \{\alpha_2\}$, $\Sigma/\Delta' = \{\alpha_1 + \alpha_2 + \alpha_3\}$ and $\mathbf{A} = \emptyset$.



In this example, the above hypothesis on the negative colors of $\mathscr{S}/\Delta'$ is not verified. One has $H \cong \mathrm{GL}(3)$ included in K, which is a parabolic subgroup of semisimple type A_2 of $\mathrm{SO}(6) \subset G$.

5.3. **The generic stabilizer of a system admitting a quotient of higher defect.**

Here we construct the generic stabilizer H of a spherical system $\mathscr{S}$ admitting a *quotient of higher defect* $\mathscr{S}/\Delta'$, i.e. whose defect is strictly greater than $\mathrm{d}(\mathscr{S})$, starting from the generic stabilzers of $\mathscr{S}/\Delta'$ and of certain localizations of $\mathscr{S}$. Notice that if Δ' is minimal and $\mathrm{d}(\mathscr{S}/\Delta') > \mathrm{d}(\mathscr{S})$ then the quotient is of type $(\mathscr{L})$.

The idea is the following. If we denote by $K \supset H$ the associated co-connected inclusion, we know that Levi parts L_K and L of resp. K and H differ only by their connected centers. If the defect of K is greater than that of H, the inclusion $Z(L_K)^\circ \supset Z(L)^\circ$ is strict. On the other hand, $Z(L_K)^\circ$ must not be contained in $\mathrm{N}_G H$; therefore H^u must not be stable under conjugation by $Z(L_K)^\circ$.

Roughly speaking, the result is that H^u is a $Z(L)^\circ$-stable (L_K°, L_K°)-submodule of K^u, diagonally embedded in a sum of L_K-submodules which are all isomorphic except for the action of $Z(L_K)^\circ$.

Moreover, these submodules can be determined inside K^u using some particular localizations of $\mathscr{S}$, having the property that they admit $\mathscr{S}/\Delta'$ as a quotient too.

More precisely, let Δ' be a distinguished subset of type $(\mathscr{L})$ with $k = \mathrm{d}(\mathscr{S}/\Delta') - \mathrm{d}(\mathscr{S})$.

Conjecture 5.3.1. *There exists a unique set of $k+1$ spherical roots $\sigma_0, \ldots, \sigma_k \in \Sigma$ such that, if we denote $\Sigma_i = \Sigma \setminus \{\sigma_0 \ldots, \sigma_{i-1}, \sigma_{i+1}, \ldots, \sigma_k\}$, $\mathscr{S}_i = \mathscr{S}_{\Sigma_i}$ and Δ'_i the restriction of Δ' to $\mathscr{S}_i$, for all $i = 0, \ldots, k$:*

1. $\mathrm{d}(\mathscr{S}_i) = \mathrm{d}(\mathscr{S}) + k$;
2. Δ'_i *is of type* $(\mathscr{L})$;
3. $\mathscr{S}_i/\Delta'_i = \mathscr{S}/\Delta'$.

Moreover, if for all $\sigma \in \Sigma$ we denote $\Delta(\sigma)$ the subset of colors D such that $c(D, \sigma) > 0$:

4. *exactly one of the colors of $\Delta(\sigma_i)$ is not free, for all $i = 0, \ldots, k$;*
5. *either $\Delta(\sigma_0) \cap \ldots \cap \Delta(\sigma_k) = \{E\}$ (with $E \in \Delta(\sigma)$ only if $\sigma = \sigma_i$ for some $0 \le i \le k$) or each $\Delta(\sigma_i)$ intersects $\Delta(\sigma)$ for some simple root $\sigma \neq \sigma_i$ for all $0 \le i \le k$.*

Finally, if we denote Δ_i the set of colors of $\mathscr{S}_i$ and c_i the Cartan pairing, for all $i = 0, \ldots, k$, there is a natural bijection $\varphi_i \colon \Delta \to \Delta_i$, such that, if Δ'' is a minimal parabolic subset of Δ containing Δ':

6. *$\varphi_i(\Delta'')$ is minimal parabolic containing Δ'_i and $\mathscr{S}_i/\varphi_i(\Delta'') = \mathscr{S}/\Delta''$,*
7. *$c(D, \sigma) = c_i(\varphi_i(D), \sigma)$ for all $D \in \Delta''$ and all $\sigma \in \Sigma_i$.*

Actually the above properties (1), (2), (3) imply all the others. We omit here the general proof: we only discuss in detail the following “not too special” case.

Proposition 5.3.1. *Suppose that $\sigma_0, \ldots, \sigma_k \in \Sigma \cap S$. Then the statements (1), (2), (3) of the above conjecture imply (4), (5), (6), (7).*

Proof. Because of (3), none of $\sigma_0, \ldots, \sigma_k$ is a spherical root of $\mathscr{S}/\Delta'$. Therefore one of the colors of $\Delta(\sigma_i)$ must be in Δ', call it D_i^+. The other color D_i^- of $\Delta(\sigma_i)$ cannot be in Δ', because in $\mathscr{S}_j$ for $j \neq i$ there would exist a color moved by the simple root σ_i, such color would be negative and contained in Δ_j, but this is impossible since Δ_j is minimal. Since σ_i is not even a summand of any spherical root of $\mathscr{S}/\Delta'$, the color D_i^- must be free. We claim that D_i^+ is not free, which proves (4).

To prove the claim, let $j \neq i$ and consider $\mathscr{S}_j$: its defect is $\mathrm{d}(\mathscr{S}) + k$ and it has k spherical roots less than $\mathscr{S}$, so the number of colors of $\mathscr{S}$ and of $\mathscr{S}_j$ must be equal. The set Δ^{2a} can be identified with Δ_j^{2a}, since the only spherical roots in $\Sigma \setminus \Sigma_j$ are simple roots. Moreover, the set $\Delta_j(\sigma_h)$ for $h \neq j$ contains one element, which can be identified with D_h^-.

It follows that Δ_j^b can be identified with $\Delta^b \cup \{D_h^- \mid h \neq j\}$. The set of the remaining colors of $\mathscr{S}$ is the following:

$$\mathbf{A}' = \mathbf{A} \setminus \{D_h^- \mid h \neq j\},$$

and the set of the remaining colors of $\mathscr{S}_j$ is $\mathbf{A}_j$. We also know from §2.5 that $\mathbf{A}_j$ is identified with a subset of $\mathbf{A}$, and we conclude that it must be $\mathbf{A}'$. Hence all colors in $\mathbf{A} \setminus \{D_h^- \mid h \neq j\}$ are moved by some simple spherical root in Σ_j.

Now Σ_j doesn't contain σ_h if $h \neq j$, hence in $\mathscr{S}$ each color D_h^+ must be moved by some spherical simple root other than σ_h. Now $j \neq i$, so D_i^+ is one of them, and our claim is proved.

We now prove (5). Suppose that there is some σ_i such that no color in $\Delta(\sigma_i)$ is moved by roots in $S' = S \setminus \{\sigma_0, \ldots, \sigma_k\}$. We must then show the existence of E, and we proceed by induction on k.

We set $E = D_i^+$ the non-free color in $\Delta(\sigma_i)$: it is moved by some simple root, not in S', so E is also equal to D_j^+ for some $j \neq i$. If $k = 1$ then $\{i, j\} = \{0, 1\}$ and we are done. Otherwise, consider the localization of $\mathscr{S}$ to $\Sigma \setminus \{\sigma_i\}$. It is not difficult to check that the restriction $\Delta'|_{\Sigma\setminus\{\sigma_i\}}$ gives a quotient of higher defect satisfying (1), (2), (3), (4), and with difference of defects equal to $k - 1$.

The colors moved by σ_j are again not moved by any root in S', so by induction $\sigma_0, \ldots, \sigma_{i-1}, \ldots, \sigma_{i+1}, \ldots, \sigma_k$ all move one color in common (in the localization). This color must be the restriction of E, and the proof of (5) is complete.

For (6) and (7) we need φ_i. We have already defined a candidate bijection φ_i in the first part of the proof, describing the identification of Δ with Δ_i needed to prove that D_i^+ is not free. For any Δ'' minimal parabolic containing Δ', we have immediately that $\varphi_i(\Delta'')$ is the restriction $(\Delta'')|_{\Sigma_i}$. From part (3) and Proposition 2.5.1 we deduce part (6).

Let us check the condition (7) on the Cartan pairing. It is obvious on all colors D moved by some simple root in S'. It is obvious also on E, if it exists: just notice that:

$$\varphi(E) = D_i^+.$$

The remaining colors are the D_i^-'s, but they do not belong to Δ'' by minimality of the latter, so we have no condition on the Cartan pairing to check (by the way, it would be false). ☐

The whole conjecture could be proved by reducing to the list of primitive spherical systems. In the following, the quotient of higher defect $\mathscr{S}/\Delta'$ will be assumed to enjoy all the above properties (these are combinatorial conditions that can be directly verified).

Suppose now that $\mathscr{S}_i = (S_i^p, \Sigma_i, \mathbf{A}_i)$ is geometrically realizable with wonderful subgroup H_i, for all $i = 0, \ldots, k$. Suppose moreover that $\mathscr{S}/\Delta'$ is geometrically realizable, with wonderful subgroup K. For all i, we can suppose that $H_i \subset K$ and choose corresponding Levi subgroups $L_i \subseteq L_K$.

The quotients $\mathscr{S}_i/\Delta_i'$ are all of type $(\mathscr{L})$; this implies that $H_i^u \subset K^u$ and that $\mathrm{Lie}\, K^u)/\mathrm{Lie}\, H_i^u$ is a simple L_i-module. Moreover L_i and L_K differ only by their connected centers, but $\mathrm{d}(\mathscr{S}_i) = \mathrm{d}(\mathscr{S}/\Delta')$, and so $L_i = L_K$ for all i. Let us denote by $Q_- = Q_-^u\, L_{Q_-}$ the parabolic subgroup (containing B_- and K) corresponding to Δ''.

We want to study the unipotent radicals H_i^u, with the aim of proving that:

$$\mathrm{Lie}\, K^u/\mathrm{Lie}\, H_i^u \cong \mathrm{Lie}\, K^u/\mathrm{Lie}\, H_j^u$$

as L_K-modules for all i, j. We obtain this isomorphism using auxiliary subgroups H_i', which differ from H_i only for the fact that the connected center of their Levi parts is smaller. A posteriori, it will be exactly the connected center of L.

We define the subgroups H_i' with the use of *augmentations* of spherical systems, see [14, §2.2] for details. From [14, §7.2] we have that a spherical subgroup $H_i' \subseteq H_i$ with $\mathrm{N}_G H_i' = \mathrm{N}_G H_i$ corresponds to the datum of a sublattice Ξ_i' of T-weights such that:

$$\mathbb{Z}\Sigma_i \subseteq \Xi_i',$$

together with a map:

$$\rho_i' \colon \mathbb{Z}\mathbf{A}_i \to \mathrm{Hom}_{\mathbb{Z}}(\Xi_i', \mathbb{Q})$$

extending the map ρ_i of $\mathscr{S}_i$, and satisfying the compatibility conditions (a1), (a2), (σ1), (σ2), (s) of [14, §2.2].

Lemma 5.3.1. *For all $i, j = 0, \ldots, k$:*

1. *The pair:*
$$(\Xi', \rho_i') = (\mathbb{Z}\Sigma, \rho),$$
where ρ is the one of $\mathscr{S}$, is an augmentation of $\mathscr{S}_i$.
2. *Call $H_i' \subseteq H_i$ the spherical subgroup associated to the augmentation $(\mathbb{Z}\Sigma, \rho)$; then:*
$$(H_i')^u = H_i^u.$$
3. *If $L_i' \subseteq L_i$ is a Levi part of H_i', then L_i' and L_i differ only by their connected centers; moreover:*
$$L_i' = L_j',$$
so we will denote this common Levi part simply by L'.
4. *The spherical L_{Q_-}-varieties:*
$$Q_-/H_i' \cong L_{Q_-} \times_{L'} \mathrm{Lie}\, Q_-^u / \mathrm{Lie}\, H_i^u$$
are all L_{Q_-}-equivariantly isomorphic.

Proof. The compatibility relations for $(\mathbb{Z}\Sigma, \rho)$ follow from the axioms of spherical system applied to $\mathscr{S}$.

Part (2) holds because $H_i \subseteq \mathrm{N}_G H_i'$, so H_i/H_i' is a diagonalizable group. The first statement of part (3) is true because $\mathbb{Z}\Sigma_i$ is a saturated sublattice of $\mathbb{Z}\Sigma$, so H_i/H_i' is connected.

We prove the second statement of part (3). Pull-back along the standard projection $G/H_i' \to G/H_i$ induces a bijection between the sets of colors of G/H_i and G/H_i', so we denote both by Δ_i. We also consider the subset $\Delta_i'' = \varphi_i(\Delta'')$ as a set of colors of both G/H_i' and G/H_i.

By hypothesis on Δ', and the fact that ρ_i' and ρ_j' are both equal to ρ, we have:

$$\rho_i'(\Delta_i'') = \rho_j'(\Delta_j'')$$

for all i, j. Moreover the colors $\Delta_i \setminus \Delta_i''$ and $\Delta_j \setminus \Delta_j''$ all have the same associated weights λ_D^* (see §3.2), since they are the weights associated to the colors of G/Q_-.

We conclude that:

$$Z(L_i')^\circ = Z(L_j')^\circ$$

for all $i, j = 0, \ldots, k$, because Lemma 3.2.1 and Corollary 3.2.1 hold for H_i' and Δ_i'' even if H_i' is not a wonderful subgroup, with exactly the same proofs.

Since L_i' and L_j' differ only by their connected centers, the equality

$$L_i' = L_j'$$

follows.

Part (4) is a consequence of the Knop conjecture, proved in [11], together with Lemma 3.3.1 (again, it holds for H'_i and Δ''_i even if H'_i is not a wonderful subgroup, with exactly the same proof).

We also point out that a much weaker version of the Knop conjecture would be enough here to prove (4), once the equality of part (3) is given. □

Corollary 5.3.1. *Up to replacing (if necessary) H_j with a conjugate also inside K and with the same Levi part L_j, we have:*

$$\operatorname{Lie} K^u / \operatorname{Lie} H_i^u \cong \operatorname{Lie} K^u / \operatorname{Lie} H_j^u$$

as irreducible L'_i-modules for all $i, j = 0, \ldots, k$.

Proof. From part (4) of Lemma 5.3.1, we deduce that the quotients $\operatorname{Lie} Q_-^u / \operatorname{Lie} H_i^u$ and $\operatorname{Lie} Q_-^u / \operatorname{Lie} H_j^u$ are L'-equivariantly isomorphic, up to twisting if necessary the action on $\operatorname{Lie} Q_-^u / \operatorname{Lie} H_j^u$ with an element $g \in \mathrm{N}_{L_{Q_-}} L'$ (see [12, Lemma 3.6.7]).

Now it is possible to apply [12, Proof of Theorem 1, step 5], with the subgroups H_1, H_2, $\widetilde{H}$ of *loc.cit.* equal resp. to H'_i, H'_j and K. The proof can be carried out in the same way even though H'_i and H'_j are not equal to their normalizers.

It follows that we may suppose $g \in \mathrm{N}_G(K)$, in other words g stabilizes K^u. Replacing H_j with gH_jg^{-1} we obtain the corollary. □

Proposition 5.3.2. *There is an L_K-module decomposition* $\operatorname{Lie} K^u = V \oplus W_0 \oplus \ldots \oplus W_k$ *and $Z(L_K)^\circ$-weights $\gamma_0, \ldots, \gamma_k$ such that:*

1. *all modules W_i are isomorphic as (L_K°, L_K°)-modules, and simple;*
2. *$Z(L_K)^\circ$ acts on W_i via the weight γ_i;*
3. $\operatorname{Lie} H_i^u = V \oplus W_0 \oplus \ldots W_{i-1} \oplus W_{i+1} \oplus \ldots \oplus W_k$.

Proof. Immediate from Lemma 5.3.1 and Corollary 5.3.1. □

Before going on, we prove the following converse to Proposition 5.3.2.

Proposition 5.3.3. *Let $\mathscr{S}$ be a geometrically realizable spherical system with generic stabilizer H and wonderful variety X, and suppose there exists a co-connected inclusion $K \supset H$, $k > 0$ and $k+1$ stabilizers $H_0, \ldots, H_k$ of points in X with the above properties. Then Δ_K satisfies the conditions of Conjecture 5.3.1.*

Proof. Let X be the wonderful embedding of G/H, and choose any $i = 0, \ldots, k$. Let λ_i be a 1-PSG of $Z(L_K)^\circ$ such that $\langle \lambda_i, \gamma_j \rangle = 0$ for $i \neq j$ and $\langle \lambda_i, \gamma_i \rangle = 1$.

Let us call $x \in X$ the point eH on the open orbit G/H. Then:

$$x^i = \lim_{t \to 0} \lambda(t)\, x$$

lies on 1-codimensional G-orbit of X for all i. We define σ_i to be the unique spherical root of X such that $\Sigma \setminus \{\sigma_i\}$ is the set of spherical roots of $\overline{Gx^i} \subset X$. With this choice, the investigated conditions follows from the properties of the co-connected inclusions $K \supset H_i$, and from Proposition 5.1.1. □

Let us come back to our previous notation and hypothesis on Δ'. We are ready to state the main result of this section, for simplicity with an additional mild assumption on K.

Recall that K is said to be *spherically closed* if all non-trivial G-equivariant automorphisms of G/K move some color. This condition is equivalent to the fact that its spherical system does not contain *loose* spherical roots of types B or G (see Definition 6.3.1).

Theorem 5.3.1. *Suppose that K is spherically closed. Then the spherical system $\mathscr{S}$ is geometrically realizable, and a generic stabilizer H can be defined as follows:*

- *its Levi part L differs from L_K only by the connected center;*
- *the connected center $Z(L)^\circ$ is the connected subgroup of $Z(L_K)^\circ$ defined by the equations $\gamma_0 = \ldots = \gamma_k$;*
- *the unipotent radical H^u is an L-submodule of K^u, in general position among those containing V.*

Proof. The subgroup H^u of K^u is well defined. We then define $L = \mathrm{N}_{L_K} H^u$: the above properties of L and $Z(L)^\circ$ follow immediately. It remains to show that H is wonderful, with spherical system $\mathscr{S}$.

Since K is spherically closed, we can use [4, §2.4.2]: for H to be wonderful it is enough that it is spherical (it is, because the subgroups H_i are spherical) and $\mathrm{N}_K H = H$ (which holds by construction).

Let X be the wonderful embedding of G/H, with spherical system $\mathscr{S}_X$: we want to show that $\mathscr{S}_X = \mathscr{S}$. We will write $\mathscr{S}_X = (S^p_X, \Sigma_X, \mathbf{A}_X)$ and $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$. Choose any $i = 0, \ldots, k$ and define λ_i be a 1-PSG of $Z(L_K)^\circ$ such that $\langle \lambda_i, \gamma_j \rangle = 1$ for $i \neq j$ and $\langle \lambda_i, \gamma_i \rangle = 0$.

This time, if $x \in X$ is the point $eH \in G/H$, then:

$$x(i) = \lim_{t \to 0} \lambda(t)\, x$$

lies on a k-codimensional orbit in X: this can be proved noticing that the projection $G/H \to G/K$ sends x to the point $eK \in G/K$, which is $\lambda_i(\mathbb{C}^*)$-stable. By construction the stabilizer $H(i)$ of $x(i)$ contains H_i and the two subgroups have the same dimension.

On the other hand, the projection $G/H \to G/K$ extends to a map between the respective wonderful embeddings, in such a way that $x(i)$ is mapped onto the point $eK \in G/K$. Hence $H(i)$ is contained in K. But the Levi subgroups L and L_K differ only by their connected centers, therefore $H(i)$ cannot contain H_i strictly: in other words $H_i = H(i)$.

What we have shown is that X has $k+1$ localizations with spherical systems $\mathscr{S}_0, \ldots, \mathscr{S}_k$. In addition, the co-connected inclusion $K \supset H$ falls under the hypotheses of Proposition 5.3.3: let Δ'_X be the corresponding distinguished subset of colors. So $\mathscr{S}$ has exactly the same quotient $\mathscr{S}/\Delta' = \mathscr{S}_X/\Delta'_X$ and same localizations $\mathscr{S}_0, \ldots, \mathscr{S}_k$: we must now prove that this implies $\mathscr{S}_X = \mathscr{S}$.

The $k+1$ localizations assure $S^p_X = S^p$ and $\Sigma_X = \Sigma$. It remains to show that $\mathbf{A}_X = \mathbf{A}$ and that the resp. bilinear pairings c_X and c are equal. It is convenient to denote by Σ' the set of common spherical roots of $\mathscr{S}_0, \ldots, \mathscr{S}_k$, i.e. $\Sigma' = \Sigma \setminus \{\sigma_0, \ldots, \sigma_k\}$, and by $\mathscr{S}'$ the localization of $\mathscr{S}$ (or of $\mathscr{S}_X$) on Σ'.

Let us choose any spherical root $\alpha \in \Sigma_X \cap S = \Sigma \cap S$ and denote by $\mathbf{A}_X(\alpha)$, $\mathbf{A}(\alpha)$ the colors moved by α in resp. $\mathscr{S}_X$, $\mathscr{S}$.

There is a bijection φ_α between $\mathbf{A}_X(\alpha)$ and $\mathbf{A}(\alpha)$ whenever $\alpha \in \Sigma'$, in such a way that $c_X(D, -) = c(\varphi_\alpha(D), -)$, where both are restricted to Σ'. This is an obvious consequence of the fact that both $\mathscr{S}$ and $\mathscr{S}_X$ localize to $\mathscr{S}'$.

This induces a bijection:

$$\varphi\colon \bigcup_{\alpha \in \Sigma'} \mathbf{A}_X(\alpha) \to \bigcup_{\alpha \in \Sigma'} \mathbf{A}(\alpha)$$

compatible with the restrictions of c_X and c on Σ'. We shall now prove that it is compatible with c_X and c on Σ, and that it extends to a bijection between $\mathbf{A}_X$ and $\mathbf{A}$.

For the first claim, let $\beta \in \Sigma \setminus \Sigma'$: then $\beta = \sigma_i$ for some i. Since both $\mathscr{S}$ and $\mathscr{S}_X$ localize to $\mathscr{S}_i$, it follows that $c_X(D, \beta) = c(\varphi(D), \beta)$.

Let us prove the second claim. First of all, suppose we have a simple root $\alpha \in \Sigma_X \setminus \Sigma'$ with a color $D \in \mathbf{A}_X(\alpha)$ moved also by a simple root in Σ'. Then, $\varphi(D)$ is already defined, with $c_X(D, -) = c(\varphi(D), -)$. Call D' the other element of $\mathbf{A}_X(\alpha)$, and set $\varphi(D')$ to be the element of $\mathbf{A}(\alpha)$ different from $\varphi(D)$. Now by the axioms of spherical systems:

$$c_X(D', -) = \langle \alpha^\vee, - \rangle - c_X(D, -) = \langle \alpha^\vee, - \rangle - c(\varphi(D), -) = c(\varphi(D'), -),$$

therefore φ respects the pairing for D' too.

We remain with the case where $\alpha \in (\Sigma \setminus \Sigma') \cap S$, and the colors in $\mathbf{A}(\alpha)$ are not moved by any simple root in Σ'. Thanks to property (5) of Conjecture 5.3.1 (which here is assumed true), all roots $\sigma_0, \ldots, \sigma_k$ are simple and move one color E in common.

This statement holds for $\mathscr{S}_X$ too, by Proposition 5.3.3. Let us denote by E_X the corresponding color. Then there is a natural bijection:

$$\varphi \colon \bigcup_{i=0}^{k} \mathbf{A}_X(\sigma_i) \to \bigcup_{i=0}^{k} \mathbf{A}(\sigma_i)$$

sending E_X to E. The identity $c_X(D, \sigma_i) = c(\varphi(D), \sigma_i)$ is thus obvious for $i = 0, \ldots, k$; the same identity for the spherical roots in Σ' follows from the localizations $\mathscr{S}_0, \ldots, \mathscr{S}_k$.

We conclude that $\mathscr{S}_X = \mathscr{S}$, and the proof is complete. □

5.4. Quotients of higher defect given by positive colors.

A particular case of quotients of higher defect has already been studied in [14] for G of type A. It is related to a special type of distinguished subsets: those containing only one element.

Definition 5.4.1. Let $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$ be a spherical system. A color D is *positive* if $c(D, \sigma) \geq 0$ for all $\sigma \in \Sigma$. In this case we define Σ_D to be the set of spherical roots σ such that $c(D, \sigma) > 0$.

It is obvious that $\Delta' = \{D\}$ is (*)-distinguished and smooth whenever D is a positive color. Indeed, $\Sigma/\{D\} = \Sigma \setminus \Sigma_D$. Since the colors of the quotient are $\Delta \setminus \{D\}$, we have:

$$\mathrm{d}(\mathscr{S}/\{D\}) - \mathrm{d}(\mathscr{S}) = |\Sigma_D| - 1.$$

Therefore $\mathrm{d}(\mathscr{S}/\{D\}) = \mathrm{d}(\mathscr{S})$ if and only if D is free.

If D is a non-free positive color of a spherical system $\mathscr{S}$, the localizations $\mathscr{S}_i$, uniquely determined by $\{D\}$ as above, contain each a free positive color equal to the restriction of D.

As a consequence, the geometric realizability of spherical systems with non-free positive colors is consequence of the geometric realizability of all spherical systems with free positive colors.

Notice that here the assumption that the generic stabilizer K of $\mathscr{S}/\{D\}$ is spherically closed can be dropped. Indeed, a loose spherical root of type B or G belongs to $\Sigma/\{D\}$ if and only if it belongs to Σ.

6. Tails

6.1. **Definition of spherical systems with tails.** Another general procedure applies to all indecomposable spherical systems $\mathscr{S}$ obtained adding one of the following spherical roots γ to another spherical system $\mathscr{S}'$ (i.e. $\mathscr{S}'$ is the localization of $\mathscr{S}$ in $S' \subset S$, with $S' \cup \operatorname{supp}\gamma = S$). We always have that γ glues on a unique connected component of S' of type A.

If $S' \cap \operatorname{supp}\gamma = \emptyset$ we denote by $\{\alpha_1, \ldots, \alpha_p\}$ (numbered from left to right) the simple roots of the connected component of S' where γ is glued. There exists at most one spherical root γ' non-orthogonal to $\operatorname{supp}\gamma$, and γ is one of the following:

b(m): if $m > 1$:



support of type B_m, $\gamma = \alpha_{p+1} + \ldots + \alpha_{p+m}$; if $m = 1$:



support of type A_1, $\gamma = \alpha_{p+1}$, with α_{p+1} the short simple root of a component of S of type B, $c(D^+_{\alpha_{p+1}}, \gamma') = c(D^-_{\alpha_{p+1}}, \gamma') = -1$.

2b(m): if $m > 1$:



support of type B_m, $\gamma = 2\alpha_{p+1} + \ldots + 2\alpha_{p+m}$; if $m = 1$:

support of type A_1, $\gamma = 2\alpha_{p+1}$, with α_{p+1} the short simple root of a component of S of type B.

d(m): if $m > 2$:



support of type D_m, $\gamma = 2\alpha_{p+1} + \ldots + 2\alpha_{p+m-2} + \alpha_{p+m-1} + \alpha_{p+m}$; if $m = 2$:



support of type $\mathsf{A}_1 \times \mathsf{A}_1$, $\gamma = \alpha_{p+1} + \alpha_{p+2}$, with $\alpha_{p+1}, \alpha_{p+2}$ simple roots at the "end" of a component of S of type D (i.e. $\{\beta \in S : \alpha_{p+1} \not\perp \beta\} = \{\beta \in S : \alpha_{p+2} \not\perp \beta\} = \{\alpha_p\}$).

If $S' \cap \operatorname{supp}\gamma \neq \emptyset$ we denote by $\{\alpha_1, \ldots, \alpha_{p+1}\}$ (numbered from left to right) the simple roots of the connected component of S' where γ is glued. Then γ is of the form:

c(m): for $m \geq 2$:



support of type C_m, $\gamma = \alpha_{p+1} + 2\alpha_{p+2} + \ldots + 2\alpha_{p+m-1} + \alpha_{p+m}$. In this case we will say that γ is *overlapping.*

The above types of tails will be called *classical.* Other cases, with S having connected components of exceptional type, can be considered as tail cases too. Here the tail $\widetilde{\Sigma} \subset \Sigma$ consists of two spherical roots. Let us denote by $\{\alpha_1, \ldots, \alpha_n\}$

the simple roots of a component of S of exceptional type, numbered as in Bourbaki. Then we define the following cases:

($\mathbf{aa},\mathbf{aa}$): component of type E_6 and $\widetilde{\Sigma}=\{\alpha_1+\alpha_6,\alpha_3+\alpha_5\}$. Example:



($\mathbf{d3},\mathbf{d3}$): component of type E_7 and $\widetilde{\Sigma}=\{\alpha_2+2\alpha_4+\alpha_5,\alpha_5+2\alpha_6+\alpha_7\}$. Example:



($\mathbf{d5},\mathbf{d5}$): component of type E_8 and $\widetilde{\Sigma}=\{2\alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\alpha_5,\alpha_2+\alpha_3+2\alpha_4+2\alpha_5+2\alpha_6\}$. Example:



($\mathbf{2a},\mathbf{2a}$): component of type F_4 and $\widetilde{\Sigma}=\{2\alpha_3,2\alpha_4\}$. Example:



Definition 6.1.1. A spherical system $\mathscr{S}$ has a *tail* in $\widetilde{\Sigma}\subset\Sigma$ if and only if there exists a (*)-distinguished subset of colors Δ'' such that $\Sigma/\Delta''=\widetilde{\Sigma}$, and: $\widetilde{\Sigma}=\{\gamma\}$ with γ one of the above cases $b(m)$, $2b(m)$, $d(m)$, $c(m)$ (in the latter case we allow $\alpha_{p+1}\in S^p/\Delta''$), or $|\widetilde{\Sigma}|=2$ and $\widetilde{\Sigma}$ is one of the above cases (aa,aa), $(d3,d3)$, $(d5,d5)$, $(2a,2a)$.

The existence of Δ'' is necessary: otherwise the generic stabilizer of $\mathscr{S}$ behaves differently from our analysis below.

A set of colors Δ'' with the property required in the above Definition always exists if $\operatorname{supp}\widetilde{\Sigma}$ is a connected component of S (so $\widetilde{\Sigma}$ is necessarily of type $c(m)$): take for example all the colors of $\mathscr{S}'$. Otherwise, if some Δ'' exists, one choice can be given by all the colors of $\mathscr{S}'$ except one. In what follows, we will suppose that Δ'' is minimal with respect to its properties.

Being of rank 1 or 2, the wonderful variety associated to $\mathscr{S}/\Delta''$ exists: we call its generic stabilizer $M\subset G$. If our strategy is based on an induction on the number of tails, we can suppose the same for the wonderful variety associated to $\mathscr{S}'$: we call its generic stabilizer $K\subset G_{S'}$. We will now see how to build a candidate subgroup H for the spherical system $\mathscr{S}$ starting from M and K.

6.2. **The generic stabilizer of a system with tails.**

6.2.1. *Tail of type $b(m)$.* An example of this kind of tail is the rank 2 spherical system of $\mathrm{SO}(2p+2m+1)$, say $p\geq 2$:



$$\mathscr{S}=(\{\alpha_2,\ldots,\alpha_{p-1},\alpha_{p+2},\ldots,\alpha_{p+m}\},\{\alpha_1+\ldots+\alpha_p,\alpha_{p+1}+\ldots+\alpha_{p+m}\},\emptyset)$$

Its wonderful subgroup has Levi factor $\mathrm{GL}(p) \times \mathrm{SO}(2m)$ and Lie algebra of the unipotent radical equal to $\mathbb{C}^p \otimes \mathbb{C}^{2m} \oplus \wedge^2\mathbb{C}^p$.

The quotient $\mathscr{S}/\Delta''$ is the rank 1 case:



whose wonderful subgroup is $M = (\mathrm{GL}(p) \times \mathrm{SO}(2m))Q^u_-$, and it is a parabolic induction by means of the parabolic subgroup $Q_- = P_{-S\setminus\{\alpha_p\}}$. The set S' is $\{\alpha_1, \ldots, \alpha_p\}$, and the localization $\mathscr{S}'$ is:



$$\mathscr{S}' = (\{\alpha_1 + \ldots + \alpha_p\}, \{\alpha_2, \ldots, \alpha_{p-1}\}, \emptyset)$$

whose wonderful subgroup is $K = \mathrm{GL}(p)$ inside $\mathrm{SL}(p+1)$. The morphism $\phi_{\Delta''}$ restricted to $\mathscr{S}'$ corresponds to the inclusion $\mathrm{GL}(p) \subset R_-$ where R_- is the parabolic subgroup of $\mathrm{SL}(p+1)$ associated to $S' \setminus \{\alpha_p\}$.

Now let us discuss a general case. We assume that $\operatorname{supp}\widetilde{\Sigma}$ is not a connected component of S, and α_p is the simple root in $S \setminus \operatorname{supp}\widetilde{\Sigma}$ with $\alpha_p \not\perp \operatorname{supp}\widetilde{\Sigma}$. Recall also that the connected component of S' containing α_p is of type A.

The homogeneous space G/M is a parabolic induction by means of the parabolic subgroup $Q_- = P_{-S''}$, minimal parabolic containing M, where $S'' = S^p/\Delta'' \cup \operatorname{supp}\widetilde{\Sigma}$. Notice that $S'' \not\ni \alpha_p$, and $G_{\operatorname{supp}\widetilde{\Sigma}}$ acts transitively on G/M: we call its stabilizer $M_{\operatorname{supp}\widetilde{\Sigma}} \subset G_{\operatorname{supp}\widetilde{\Sigma}}$. In other words, $G_{S''} = G_{S'\cap S''} \times G_{\operatorname{supp}\widetilde{\Sigma}}$ and $M = (G_{S'\cap S''} \times M_{\operatorname{supp}\widetilde{\Sigma}})Q^r_-$.

Here, in case of tail of type $b(m)$, we have $M_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{SO}(2m)$ included in $G_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{SO}(2m+1)$.

On the other hand let us restrict the set of colors Δ'' to S', and call it $\Delta''|_{S'}$. From Definition 6.1.1 and the assumption that Δ'' is minimal, it follows that $\Delta''|_{S'}$ is a minimal parabolic set of colors of $\mathscr{S}'$. The corresponding parabolic subgroup R_- of $G_{S'}$ has $S' \cap S''$ as subset of associated simple roots. Now choose K contained in R_-, which implies $K^u \subseteq R^u_-$, and choose a Levi part L_K of K contained in L_{R_-}.

By construction there exists an isogeny $\pi\colon G_{S'\cap S''}Z(L_{S''}) \to L_{R_-}$, we can set $L = \pi^{-1}(L_K) \times M_{\operatorname{supp}\widetilde{\Sigma}}$ as a subgroup of $L_{S''}$. Moreover, the L_K-submodule K^u of R^u_- corresponds to a L-stable subalgebra $\operatorname{Lie} U$ of $\operatorname{Lie} Q^u_-$, in such a way that the two quotients $\operatorname{Lie} R^u_-/\operatorname{Lie} K^u$, $\operatorname{Lie} Q^u_-/\operatorname{Lie} U$ are isomorphic as $\pi^{-1}(L_K)$-modules.

To describe more explicitly this isomorphism, let us denote with $\tilde{R}_-$ the parabolic subgroup of $G_{S'}$ associated to the subset of simple roots $S' \setminus \{\alpha_p\}$, and with $\tilde{Q}_-$ the parabolic subgroup of G associated to $S \setminus \{\alpha_p\}$.

The $L_{\tilde{R}_-}$-module $\operatorname{Lie} \tilde{R}^u_-$ is simple of highest weight $-\alpha_p$, that is, the fundamental weight $\omega_{\alpha_{p-1}}$ once restricted to the maximal torus of $G_{S'\setminus\{\alpha_p\}}$. On the other hand the quotient $L_{\tilde{Q}_-}$-module $\operatorname{Lie} \tilde{Q}^u_-/[\operatorname{Lie} \tilde{Q}^u_-, \operatorname{Lie} \tilde{Q}^u_-]$ is simple of highest weight $-\alpha_p$, that is, $\omega_{\alpha_{p-1}} + \omega_{\alpha_{p+1}}$ once restricted to the maximal torus of $G_{S\setminus\{\alpha_p\}}$. Therefore, $\operatorname{Lie} \tilde{Q}^u_-/[\operatorname{Lie} \tilde{Q}^u_-, \operatorname{Lie} \tilde{Q}^u_-] \cong \operatorname{Lie} \tilde{R}^u_- \otimes V(\omega_{\alpha_{p+1}})$, where $V(\omega_{\alpha_{p+1}})$ is the simple $\mathrm{SO}(2m+1)$-module of highest weight $\omega_{\alpha_{p+1}}$. The module $V(\omega_{\alpha_{p+1}})$ decomposes into $W(\omega_{\alpha_{p+1}}) \oplus W(0)$ as $\mathrm{SO}(2m)$-module. We obtain the above isomorphism between the quotients $\operatorname{Lie} R^u_-/\operatorname{Lie} K^u$ and $\operatorname{Lie} Q^u_-/\operatorname{Lie} U$ since $[\operatorname{Lie} \tilde{Q}^u_-, \operatorname{Lie} \tilde{Q}^u_-]$ and $\operatorname{Lie} \tilde{R}^u_- \otimes W(\omega_{\alpha_{p+1}})$ are included in $\operatorname{Lie} U$, and $\operatorname{Lie} \tilde{R}^u_-$ is isomorphic to $\operatorname{Lie} \tilde{R}^u_- \otimes W(0) \subset \operatorname{Lie} \tilde{R}^u_- \otimes V(\omega_{\alpha_{p+1}})$.

The candidate H is then $H = U\,L$.

6.2.2. *Tail of type* $2b(m)$. This case is similar to the preceeding, the only difference is that here $M_{\mathrm{supp}\,\widetilde{\Sigma}} = \mathrm{N}_{\mathrm{SO}(2m+1)}\mathrm{SO}(2m)$.

6.2.3. *Tail of type* $d(m)$. This case is similar to the preceeding: here $M_{\mathrm{supp}\,\widetilde{\Sigma}} = \mathrm{SO}(2m-1)$ inside $G_{\mathrm{supp}\,\widetilde{\Sigma}} = \mathrm{SO}(2m)$.

6.2.4. *Tail of type* $c(m)$. This case is special: the idea is similar but there are some complications.

A generic stabilizer for $\mathscr{S}'$ can be contained in two non-conjugated parabolic subgroups that are minimal containing it. For other types of tails, whenever a given $\mathscr{S}'$ appears, always the same parabolic subgroup plays the role of R_- as above. With this kind of tails different R_- appear, and they give rise to different spherical systems $\mathscr{S}$ with same tail and same localization $\mathscr{S}'$.

For a given $\mathscr{S}$, the right choice of R_- is assured by the parabolic subset $\Delta''|_{S'}$. Let us see an example based on the spherical system $\mathscr{S}'$:



Its generic stabilizer can be chosen to have Levi subgroup equal to the standard Levi of the parabolic $P_{-\{\alpha_1\}}$ of SL(4), or of the parabolic $P_{-\{\alpha_3\}}$ (see [5, Table 3, Case 9 with $p = 1$]). Both $P_{-\{\alpha_1\}}$ and $P_{-\{\alpha_3\}}$ are valid choices for R_-, and appear if $\mathscr{S}$ is resp. one of the following two tail cases:





Even when the same parabolic subgroup R_- occurs, different choices of K inside R_- may produce non-conjugated subgroups H.

Lemma 6.2.1. *Suppose that* $\mathscr{S}$ *is a tail spherical system with* $\mathscr{S}'$ *geometrically realizable and tail* $\widetilde{\Sigma}$. *Then there exists a subset of colors* $\Delta' \subseteq \Delta''$ *of* $\mathscr{S}$ *of type* $(\overline{\mathscr{L}})$ *such that* $(\Delta'|_{S'}, \Delta''|_{S'})$ *is of type* $(\overline{\mathscr{L}}, \overline{\mathscr{R}})$.

Proof. Since we assumed that Δ'' is minimal with respect to its properties, the localization $\Delta''|_{S'}$ is minimal parabolic. It contains then a subset of colors Δ_0 which has the required properties of $\Delta'|_{S'}$.

Now the colors of $\mathscr{S}'$ are identified with a subset of the colors of $\mathscr{S}$, and it is immediate to prove that we can take Δ' to be Δ_0 seen as a subset of colors of $\mathscr{S}$. ☐

Any Δ' with the above properties will determine the right choice for K. Let us see an example, $\mathscr{S}'$:



Here two subsets of type $(\overline{\mathscr{L}})$ exist, giving as $\mathscr{S}'/(\Delta'|_{S'})$ resp. the following quotients:





The two subsets of colors appear as $\Delta'|_{S'}$ if $\mathscr{S}$ is resp. one of the following two tail cases:



Let us now define the candidate subgroup H in general. Denote by Q_- the parabolic subgroup of G defined by the minimal parabolic subset containing Δ'', and by R_- the parabolic subgroups of $G_{S'}$ defined by $\Delta''|_{S'}$. Notice that α_{p+1} belongs to their sets of associated simple roots if and only if $\alpha_{p+1} \in S^p/\Delta''$.

For convenience of notation, let us also define parabolic subgroups Q'_- and R'_- obtained resp. from Q_- and R_- by adding α_{p+1} to their sets of associated simple roots. Of course, α_{p+1} may already be there, thus giving $Q'_- = Q_-$ and $R'_- = R_-$. In any case, $Q'_- = P_{-S''}$ where as above $S'' = S^p/\Delta'' \cup \operatorname{supp}\gamma$.

We have $G_{S''} = G_{S'\cap S^p/\Delta''} \times \mathrm{Sp}(2m)$, and $G_{S'\cap S''} = G_{S'\cap S^p/\Delta''} \times \mathrm{SL}(2)$. We choose a subgroup $\mathrm{SL}(2)\times\mathrm{Sp}(2m-2)$ inside $\mathrm{Sp}(2m) \subseteq G_{S''}$, and identify its $\mathrm{SL}(2)$ factor with the one of $G_{S'\cap S''}$.

Now we choose K inside R_-, in such a way that $K \subseteq R_-$ factors through the co-connected inclusion associated to $\Delta'|_{S'}$. Again we have $K^u \subsetneq R_-^u$, and we choose a Levi component L_K of K contained in L_{R_-}.

As before, we set $L = \pi^{-1}(L_K)\times\mathrm{Sp}(2m-2)$, and define U in the same way: notice that the simple $\mathrm{Sp}(2m)$-module $V(\omega_{\alpha_{p+1}})$ decomposes into $W_1 \oplus W_2$ as $\mathrm{SL}(2)\times\mathrm{Sp}(2m-2)$-module, where W_1 and W_2 are the defining modules resp. of the first and second factor. Finally, put $H = U\,L$.

From the proof of Lemma 6.3.3 it will be evident why in the other cases the definition of H does not require the use of Δ'.

6.2.5. *Tail of type* (aa, aa). The case is similar to the above classical cases $b(m)$, $2b(m)$ and $d(m)$: here $M_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{N}(\mathrm{diag}(\mathrm{SL}(3)))$ inside $G_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{SL}(3)\times\mathrm{SL}(3)$.

6.2.6. *Tail of type* $(d3, d3)$. The case is similar to the preceeding: here $M_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{N}(\mathrm{Sp}(6))$ inside $G_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{SL}(6)$.

6.2.7. *Tail of type* $(d5, d5)$. The case is similar to the preceeding: here $M_{\operatorname{supp}\widetilde{\Sigma}}$ is the normalizer of a simple group of type F_4 inside $G_{\operatorname{supp}\widetilde{\Sigma}}$ simple of type E_6.

6.2.8. *Tail of type* $(2a, 2a)$. The case is similar to the preceeding: here $M_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{N}(\mathrm{SO}(3))$ inside $G_{\operatorname{supp}\widetilde{\Sigma}} = \mathrm{SL}(3)$.

6.3. **Geometric realizability of a system with tails.** We use here all the elements previously introduced. Denote also by M' the factor $\mathrm{Sp}(2m-2)$ of L acting trivially on Q_-^u/U if the tail is of type $c(m)$, or the factor $M_{\operatorname{supp}\widetilde{\Sigma}}$ previously defined otherwise.

From now on, let $\mathscr{S}$ be a spherical system with tail in $\widetilde{\Sigma}$ and let H be the subgroup defined as above. Moreover, suppose that H is a generic stabilizer of a wonderful variety X with spherical system $\mathscr{S}_X$.

Lemma 6.3.1. *The set $\widetilde{\Sigma}$ is included in Σ_X. In the case of a tail $\widetilde{\Sigma} = \{\gamma\}$ of type $b(1)$, both colors moved by γ are free, and the Cartan pairing is equal on each of them.*

Proof. Let us choose a 1-PSG $\lambda\colon \mathbb{C}^* \to T$ adapted to $\operatorname{supp}\widetilde{\Sigma}$ (see [13]), and consider the definition of the candidate subgroup H. It is not B-spherical, in other words it is the stabilizer of a point $x \in X$ not lying on the open B-orbit. Nevertheless, uHu^{-1} is B-spherical for any u varying in a non-empty open subset of B^u_-. This means that ux lies on the open B-orbit of X, and thus the limit:

$$x_0 = \lim_{t\to 0} \lambda(t)ux$$

lies on the open $G_{\operatorname{supp}\widetilde{\Sigma}}$-orbit of the localization $X_{\operatorname{supp}\widetilde{\Sigma}}$ of X in $\operatorname{supp}\widetilde{\Sigma}$ (see [13, §1.1]). It is possible to write any $u \in B^u_-$ as a product $u = vw$ such that $w \in H$ and v commutes with M'. The consequence is that $M' \subset uHu^{-1}$ for any $u \in B^u_-$. The elements $\lambda(t)$ commute with M' for any t: we conclude that M' stabilizes x_0, and hence it is contained in a generic stabilizer of $X_{\operatorname{supp}\widetilde{\Sigma}}$. Now it is easy to list all spherical subgroups of $G_{\operatorname{supp}\widetilde{\Sigma}}$ containing M': they have rank 1 or 2 and in each case $\widetilde{\Sigma} \subset \Sigma_X$. If the tail $\widetilde{\Sigma} = \{\gamma\}$ is of type $b(m)$ one actually has that γ or 2γ is a spherical root: the case 2γ is excluded by considering the group of G-equivariant automorphisms of G/H.

For the case $b(1)$, the fact that the two colors are free stems from the inclusion $H \subseteq M$, which gives rise to a quotient of $\mathscr{S}_X$ where the assertion is true. The equality of the Cartan pairing on the two colors is easily deduced using the same tail system where γ is replaced by 2γ, and comparing the two candidate subgroups. □

Corollary 6.3.1.

(1) *Any spherical root $\sigma \in \Sigma_X \setminus \widetilde{\Sigma}$ satisfies $\operatorname{supp}\sigma \subseteq S'$.*
(2) *The spherical system $\mathscr{S}_X$ has a tail in $\widetilde{\Sigma}$.*

Proof. The first assertion is immediate from the shape of the tail spherical roots: it is enough to start from $\widetilde{\Sigma}$ and then try to add another spherical root to it using the well known classification of wonderful varieties of rank 2.

For the second assertion we must verify Definition 6.1.1 and show the existence of a (*)-distinguished subset of colors Δ''_X: this is just the subset of colors associated to the inclusion $H \subseteq M$. □

Lemma 6.3.2. *The localization of $\mathscr{S}_X$ in S' is equal to $\mathscr{S}'$.*

Proof. We suppose at first that the tail is not of type $b(1)$, and let us call H' a generic stabilizer of $X_{S'}$. We claim that H' can be chosen to have L_K as Levi factor.

Corollary 6.3.1 implies that there is a bijection between the set of distinguished sets of colors of X contained in Δ''_X, and the set of distinguished sets of colors of $X_{S'}$ contained in $\Delta''_X|_{S'}$. The bijection is simply induced by restriction, which in this case is just considering a set of colors of X as a set of colors of $X_{S'}$.

This bijection preserves the property of being of type $(\overline{\mathscr{L}})$. On the other hand $M_{\operatorname{supp}\widetilde{\Sigma}}$ is a very reductive subgroup of $L_{\operatorname{supp}\widetilde{\Sigma}}$ (i.e. not contained in any proper parabolic): therefore there exists a subset of colors Δ'_X of type $(\overline{\mathscr{L}})$ inside Δ''_X. It follows that H' has Levi factor which differs from L_K only for its connected center.

Let $\Delta'''_X \supseteq \Delta''_X$ be a minimal parabolic subset of colors of $\mathscr{S}_X$. From Lemma 6.3.1 we deduce that Δ'''_X contains all colors moved by simple roots in $S \setminus S'$, and that $\Delta'''_X|_{S'}$ is minimal parabolic of $\mathscr{S}|_{S'}$. Moreover, on $\widetilde{\Sigma}$ all colors in Δ''_X are 0 and all colors in $\Delta_X \setminus \Delta'''_X$ are ≤ 0.

From these facts it is easy to deduce that the formula of Corollary 3.2.1 gives the same result whether applied to X and Δ'''_X or to $X_{S'}$ and $\Delta'''_X|_{S'}$: for each type of tail it is an elementary exercise involving the values on $\widetilde{\Sigma}$ of the color(s) moved by the simple root(s) in $S \setminus S'$. Comparing the connected center of L and of L_K we conclude that L_K and a Levi subgroup of H' also have same connected center, and our claim follows.

Now we prove that H' and K are conjugated. It is possible to choose a 1-PSG $\lambda\colon \mathbb{C}^* \to T$ such that $\lambda(t)$ commutes with $L_{S'}$ for all $t \in \mathbb{C}^*$, and $\langle \lambda, \alpha \rangle = 1$ for all $\alpha \in S \setminus S'$. Corollary 6.3.1 implies that $\langle \lambda, \sigma \rangle > 0$ for all $\sigma \in \widetilde{\Sigma}$, and $\langle \lambda, \sigma \rangle = 0$ for all $\sigma \in \Sigma_X \setminus \widetilde{\Sigma}$. The consequence is that the limit:

$$x_0 = \lim_{t \to 0} \lambda(t) x$$

lies on the open G-orbit of $X_{\Sigma_X \setminus \widetilde{\Sigma}}$, and $\pi^{-1}(L_K)$ is contained in the stabilizer $\lambda(t) H \lambda(t)^{-1}$ of $\lambda(t)x$ for all $t \in \mathbb{C}^*$. Hence it is also contained in the stabilizer $H_{\Sigma \setminus \widetilde{\Sigma}}$ of x_0.

Thanks to the definition of H, for a generic choice of $M_{\mathrm{supp}\,\widetilde{\Sigma}}$ inside $L_{\mathrm{supp}\,\widetilde{\Sigma}}$ we have:

$$\lim_{t \to 0} \mathrm{Ad}(\lambda(t)) \left(\mathrm{Lie}\, L_K \oplus \mathrm{Lie}\, U \right) = \mathrm{Lie}\, L_K \oplus \mathrm{Lie}\, K^u.$$

where we identify $\mathrm{Lie}\, G_{S' \cap S''} \oplus \mathrm{Lie}\, Z(L_{S''})$ with $\mathrm{Lie}\, L_{R_-}$ via the isogeny π.

Hence $\mathrm{Lie}\, L_K \oplus \mathrm{Lie}\, K^u$ is also contained in $\mathrm{Lie}\, H_{\Sigma \setminus \widetilde{\Sigma}}$. Finally, Lemma 6.3.1 and Corollary 6.3.1 assure that $X_{\Sigma \setminus \widetilde{\Sigma}}$ is the parabolic induction of $X_{S'}$ by means of a parabolic P containing $P_{-S'}$. In other words $\mathrm{Lie}\, H_{\Sigma \setminus \widetilde{\Sigma}}$ is the sum of $\mathrm{Lie}\, H'$ and the radical of $\mathrm{Lie}\, P$, up to conjugating H' if necessary. We deduce that $H' \supseteq L_K\, K^u = K$, and for dimension reasons they are equal.

If the tail $\widetilde{\Sigma} = \{\gamma\}$ has type $b(1)$ we can apply Lemma 6.3.1, Corollary 6.3.1 and the first part of this proof to the spherical system $\mathscr{T}$ obtain adding to $\mathscr{S}'$ the tail $\{2\gamma\}$. We only need to check that the candidate subgroup N of $\mathscr{T}$ is wonderful too, but this stems from the fact that $H \subseteq N \subseteq \mathrm{N}_G H$.

Let Y be the wonderful completion of G/N, and $\mathscr{S}_Y$ its spherical system. The first part of the proof assures that the localization of $\mathscr{S}_Y$ on S' is equal to $\mathscr{S}'$. On the other hand $[N : H] = 2$ together with $\gamma \in \Sigma_X$, $2\gamma \in \Sigma_Y$ imply that the localizations of $\mathscr{S}_X$ and of $\mathscr{S}_Y$ in S' are equal: the proof is complete. □

Lemma 6.3.3. *The system $\mathscr{S}_X$ is equal to $\mathscr{S}$.*

Proof. Applying Lemma 6.3.1 and Lemma 6.3.2 we know that $\widetilde{\Sigma} \subset \Sigma_X$ and that the localization of $\mathscr{S}_X$ in S' is $\mathscr{S}'$. This finishes the proof if the tail is not of type $c(m)$. If $\widetilde{\Sigma} = \{\gamma\}$ is of type $c(m)$, the systems $\mathscr{S}_X$ and $\mathscr{S}$ may both share these properties, and yet be different. Namely, there can be two different choices for the color D of $\mathscr{S}'$ such that $c(D, \gamma) < 0$. If this is the case, these colors are both moved by α_p (in the notation of §6.2.4), they are both free, and both choices lead to consistent spherical systems. We must now show that one of the two possible spherical systems does not satisfy Definition 6.1.1, so there is no ambiguity, or that for each of the two systems $\mathscr{S}$ our procedure provides the right generic stabilizer H.

It is convenient here to denote by D_n^+ and D_n^- the two colors moved by a spherical root $\alpha_n \in S \cap \Sigma$.

Step 1. Let us suppose that no spherical root $\sigma \neq \gamma$ is supported on α_{p+1}. This implies that $\mathscr{S}'$ must be non-cuspidal, with α_{p+1} outside the support of its spherical roots. The system $\mathscr{S}$ in the vicinity of γ is as follows:



or:



depending on the choice of D. If D_{p+2} is the unique color moved by α_{p+2}, then Δ'' and $\{D_{p+2}\}$ decompose $\mathscr{S}$ in both cases, therefore it is not a tail system.

As a consequence, we may assume that some spherical root $\sigma \neq \gamma$ is supported on α_{p+1}.

Step 2. Let's consider first the case where $\sigma \notin S$. The only possibility is $\gamma = \alpha_{p+1} + \alpha_{p-1}$, and $\mathscr{S}$ in the vicinity of γ looks like:



or:



The required set of colors Δ'' exists only in the second diagram, so there is no ambiguity.

Step 3. It remains the case where $\alpha_{p+1} \in \Sigma$. If D_{p+1}^+ and D_{p+1}^- are free, then $\mathscr{S}$ in the vicinity of γ is:



or:



The two quotients $\mathscr{S}/\Delta'$ are resp.:





and they give two different candidates for H. The existence of the co-connected inclusions associated to the two quotients above assures that the two subgroups H correspond to the right spherical systems.

We may now assume that D_{p+1}^+ and D_{p+1}^- are not both free.

Step 4. Let us suppose that both are non-free: one of them must take the value -1 on α_p, say D^+_{p+1}. Consider the colors D^+_p and D^-_p moved by α_p. If one of them is zero both on α_{p-1} and on α_{p+1}, then we have:



or:



The first system does not admit a subset of colors Δ'', so there is no ambiguity.

We assume now that no color moved by α_p is zero on both α_{p-1} and α_{p+1}. If D^-_{p-1} is free, then we have:



or:



Here the two quotients $\mathscr{S}/\Delta'$ must be resp.:



and:



and they give two different candidates for H. Again, the existence of the co-connected inclusions associated to the two quotients above assures that the two subgroups H correspond to the right spherical systems.

If D^-_{p-1} is not free, then we have:



The two quotients $\mathscr{S}/\Delta'$ are resp.:





and they give two different candidates for H. Again, the existence of the co-connected inclusions associated to the two quotients above assures that the two subgroups H correspond to the right spherical systems.

Step 5. Let us suppose that only one of D^+_{p+1}, D^-_{p+1} is not free, say D^+_{p+1}. We leave for a moment undetermined the values of the colors D^+_p and D^-_p, and focus on the value of D^+_{p+1} on α_p. We have a system like:



if $c(D^+_{p+1}, \alpha_p) = 0$, or:



if $c(D^+_{p+1}, \alpha_p) = -1$. Since D^-_{p+1} is free, the color E must also be free in both cases.

Hence in the first case D^+_{p+1} is a non-free positive color, and the system admits a good quotient of higher defect. Thus, we may assume that $c(D^+_{p+1}, \alpha_p) = -1$.

Step 6. Now let us consider the colors moved by α_p. If one of them is zero both on α_{p-1} and on α_{p+1}, then we have:



or:



The first of the two systems does not admit a subset of colors Δ'', so there is no ambiguity.

Step 7. We may now assume that no color moved by α_p is zero both on α_{p-1} and α_{p+1}. Our system is:



or:



Both are tail systems. Since D^-_{p+1}, D^+_p, D^-_p are free, we conclude that D^-_{p-1} is free, and that D^+_{p-1}, D^-_{p-1}, D^+_{p-1} are all zero on spherical roots different from α_{p-1}, α_p, α_{p+1}, γ.

It is easy to deduce that the two quotients by Δ'' are resp.:



and:



It is evident that these correspond to two different parabolic subgroups R_-, which means two candidate wonderful subgroups. Finally, the existence of the co-connected inclusions associated to the two quotients above assures the good choice of our generic stabilizers. $\square$

Before stating the main theorem of this section, we need the following auxiliary result which is independent from the rest of the paper.

Definition 6.3.1. [4, §2.2] Let σ be a spherical root of a spherical system $\mathscr{S}$. Then σ is *loose* if and only if one of the following holds:

(A) $\sigma \in \Sigma \cap S$ and $\rho(D^+) = \rho(D^-)$ where $\Delta(\alpha) = \{D^+, D^-\}$;

(B) $\sigma = \alpha_1 + \ldots + \alpha_m$ where $\{\alpha_1, \ldots, \alpha_m\} \subseteq S$ has type B_m with short root α_m, and $\Delta(\alpha_i) = \varnothing$ for $i = 2, \ldots, m$;

(G) $\sigma = 2\alpha_1 + \alpha_2$ where $\{\alpha_1, \alpha_2\} \subseteq S$ has type G_2 with short root α_1.

Lemma 6.3.4. *Let $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$ be a spherical system. Let $\overline{\Sigma}$ be defined as Σ where we replace each loose spherical root σ with 2σ, and let $\overline{\mathbf{A}}$ be obtained from $\mathbf{A}$ by removing all pairs $\mathbf{A}(\alpha)$ for $\alpha \in \Sigma \cap S$ loose of type* A. *Then $\mathrm{N}_G(\mathscr{S}) = (S^p, \overline{\Sigma}, \overline{\mathbf{A}})$ is a spherical system; if is geometrically realizable, then $\mathscr{S}$ is geometrically realizable too.*

Proof. The lemma follows from [12, Theorem 2] and the theory of augmentations of spherical systems of [14]. We use here an alternative definition of spherical roots, valid for any spherical homogeneous space, and given in [12].

The fact that $\mathrm{N}_G(\mathscr{S})$ is a spherical system follows directly from the definition of $\overline{\Sigma}$. Suppose that $\mathrm{N}_G(\mathscr{S})$ is geometrically realizable, with generic stabilizer N: from [12, Theorem 2] we see that N is equal to its normalizer in G.

The lattice $\mathbb{Z}\Sigma \supseteq \mathbb{Z}\overline{\Sigma}$ and the Cartan pairing of $\mathscr{S}$ define an augmentation of $\mathrm{N}_G(\mathscr{S})$ in the sense of [14]. Let $H \subseteq N$ be the corresponding spherical subgroup. Using the same approach as [9, §4] we can compare the cones of G-invariant discrete valuations of G/H and of $G/\overline{H}$: since $[\mathbb{Z}\overline{\Sigma} : \mathbb{Z}\Sigma]$ is finite, we deduce that we can identify $\mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}\Sigma, \mathbb{Q})$ and $\mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}\overline{\Sigma}, \mathbb{Q})$ in such a way that the cones of invariant valuations are equal to the same subset V, a co-simplicial strictly convex polyhedral cone.

We have that V is defined by the equations $\langle \cdot, \sigma \rangle \leq 0$ for all $\sigma \in \overline{\Sigma}$. But $|\Sigma| = |\overline{\Sigma}|$ and for all $\sigma \in \overline{\Sigma}$ either σ or $\sigma/2$ is in Σ, therefore V is also defined by the equations $\langle \cdot, \sigma \rangle \leq 0$ for all $\sigma \in \Sigma$.

It follows that Σ is by definition the set of spherical roots of G/H. Since it generates $\mathbb{Z}\Sigma$ and the latter is the lattice of B-weights of $\mathbb{C}(G/H)$, it follows that H is wonderful with spherical system $\mathscr{S}$. □

Theorem 6.3.1. *Let $\mathscr{S}$ be a spherical system with a tail, and suppose that the localization $\mathscr{S}'$ as defined in §6.1 is geometrically realizable. Then $\mathscr{S}$ is geometrically realizable, with generic stabilizer H as defined in §6.2.*

Proof. We begin by supposing that the tail is not of type $b(m)$, and that $\mathscr{S}'$ corresponds to a wonderful variety without non-trivial G-equivariant automorphisms. Then $\mathrm{N}_{L_{S'}} K = K$, and from the definition of H it follows that $\mathrm{N}_G H = H$. The subgroup H is hence wonderful, call X its wonderful variety. Applying Lemma 6.3.1, Lemma 6.3.2 and Lemma 6.3.3 we know that $\mathscr{S}_X = \mathscr{S}$.

Now drop our assumptions both on $\mathscr{S}'$ and on the tail. Let $\mathrm{N}_G(\mathscr{S})$ be defined as in Lemma 6.3.4: it is a tail spherical system with localization $\mathrm{N}_G(\mathscr{S}')$ and same tail as $\mathscr{S}$, except for the case where the latter has tail $b(m)$ and $\mathrm{N}_G(\mathscr{S})$ has tail $2b(m)$. By the first part of the proof $\mathrm{N}_G(\mathscr{S})$ is geometrically realizable (with wonderful subgroup N), and therefore $\mathscr{S}$ too, thanks to Lemma 6.3.4. Let $\widetilde{H}$ be the generic stabilizer of $\mathscr{S}$: it remains to show that $\widetilde{H} = H$.

The group $\widetilde{H}$ is obtained from N in the proof of Lemma 6.3.4 by means of the augmentation of $\mathrm{N}_G(\mathscr{S})$ given by $\mathscr{S}$. Hence $\widetilde{H}$ is defined inside N as the intersection of the kernels of certain characters, we refer to [14, §6] for details. The same considerations apply to the generic stabilizers K and M of resp. $\mathscr{S}'$ and $\mathrm{N}_G(\mathscr{S}')$. Moreover, H has (by its definition) finite index in N, so it is the intersection of the kernels of some characters too.

At this point the only thing to check is that $\widetilde{H}$ and H have Levi subgroups equal up to conjugation. Let (Δ', Δ'') be as in Lemma 6.2.1, and let $\Delta''' \supseteq \Delta''$ be a minimal parabolic subset of colors. Using Proposition 3.1.1 and Corollary 3.2.1 we conclude that a Levi subgroup of $\widetilde{H}$ has the form $\pi^{-1}(L_K) \times M_{\mathrm{supp}\,\widetilde{\Sigma}}$, with the notation of §6.2. The equality $\widetilde{H} = H$ follows. □

6.4. **The generic stabilizer of a system with arcs of type $a(m)$.** Let us now mention a variant of the above situation where the same arguments apply.

Let $\mathscr{S}$ be a spherical system such that

- there exists $\gamma \in \Sigma$ of type $a(m)$,
- there exist $\gamma', \gamma'' \in \Sigma$ non-orthogonal to $\mathrm{supp}\,\gamma$ with $\mathrm{supp}\,\gamma' \cap \mathrm{supp}\,\gamma'' = \emptyset$ and such that, if $\{D \in \Delta : c(D,\gamma) = 1\} = \{D', D''\}$, then $c(D', \gamma') = c(D'', \gamma'') = -1$ and $c(D', \gamma'') = c(D'', \gamma') = 0$,
- there exists a (*)-distinguished subset of colors Δ'' such that $\Sigma/\Delta'' = \{\gamma\}$.

In this case we say that $\mathscr{S}$ has an *arc* in $\{\gamma\}$ of type $a(m)$.

Example:



Notice the analogies with the tail cases: $\mathscr{S}$ is obtained adding γ to a spherical system $\mathscr{S}'$ (localization of $\mathscr{S}$ in $S' = S \setminus \mathrm{supp}\,\gamma$). Let us enumerate the simple roots of the connected component of S containing $\mathrm{supp}\,\gamma$ in such a way that $\gamma = \alpha_{p+1} + \ldots + \alpha_{p+m}$. Actually, if $\mathscr{S}$ is indecomposable, without positive n-combs with $n > 1$, the only two simple roots of S' non-orthogonal to $\mathrm{supp}\,\gamma$ are α_p and α_{p+m+1}. As above, Δ'' is supposed to be minimal with respect to its properties, $M \subset G$ and $K \subset G_{S'}$ are resp. the generic stabilizers of $\mathscr{S}/\Delta''$ and $\mathscr{S}'$.

The description of the subgroup H and the proof that it is actually the generic stabilizer of $\mathscr{S}$ can be conducted as above. Here, take $Q_- = P_{-S''}$ with $S'' = S^p/\Delta'' \cup \mathrm{supp}\,\gamma$ and notice that $S'' \not\ni \alpha_p, \alpha_{p+m+1}$. Here we have $M_{\mathrm{supp}\,\gamma} = \mathrm{GL}(m)$ included in $G_{\mathrm{supp}\,\gamma} = \mathrm{SL}(m{+}1)$. Take R_-, the parabolic subgroup of $G_{S'}$ associated to $S' \cap S''$. Choose $K \subset R_-$ (thus $K^u \subset R^u_-$) and $L_K \subset L_{R_-}$. As above, set $L = \pi^{-1}(L_K) \times M_{\mathrm{supp}\,\gamma} \subset L_{S''}$ and $U \subset Q^u_-$ such that $\mathrm{Lie}\,R^u_-/\mathrm{Lie}\,K^u$, $\mathrm{Lie}\,Q^u_-/\mathrm{Lie}\,U$ are isomorphic as $\pi^{-1}(L_K)$-modules.

References

[1] P. Bravi, *Wonderful varieties of type E*, Represent. Theory **11** (2007) 174–191.

[2] P. Bravi, *Primitive spherical systems*, arXiv:0909.3765v2 .

[3] P. Bravi, S. Cupit-Foutou, *Classification of strict wonderful varieties*, Ann. Inst. Fourier (Grenoble) **60** (2010) 641–681.

[4] P. Bravi, D. Luna, *An introduction to wonderful varieties with many examples of type F4*, J. Algebra **329** (2011) 4–51.

[5] P. Bravi, G. Pezzini, *Wonderful varieties of type D*, Represent. Theory **9** (2005) 578–637.

[6] M. Brion, *Classification des espaces homogènes sphériques*, Compos. Math. **63** (1987), 189–208.

[7] S. Cupit-Foutou, *Invariant Hilbert schemes and wonderful varieties*, arXiv:0811.1567v2 .

[8] S. Cupit-Foutou, *Wonderful varieties: a geometrical realization*, arXiv:0907.2852v3 .

[9] F. Knop, *Automorphisms, root systems, and compactifications of homogeneous varieties*, J. Amer. Math. Soc. **9** (1996), no. 1, 153–174.

[10] F. Knop, B. Van Steirteghem, *Classification of smooth affine spherical varieties*, Transform. Groups **11** (2006), 495–516.

[11] I.V. Losev, *Proof of the Knop conjecture*, Ann. Inst. Fourier (Grenoble) **59** (2009) 1105–1134.

[12] I.V. Losev, *Uniqueness property for spherical homogeneous spaces*, Duke Math. J. **147** (2009) 315–343.

[13] D. Luna, *Grosses cellules pour les variétés sphériques*, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., 9, Cambridge Univ. Press, Cambridge, 1997, 267–280.

[14] D. Luna, *Variété sphériques de type A*, Publ. Math. Inst. Hautes Études Sci. **94** (2001) 161–226.

[15] G. Pezzini, *Wonderful varieties of type C*, Ph.D. Thesis, Università La Sapienza, Rome, 2004.